arXiv:1502.05446v1 [nlin.CD] 19 Feb 2015

Chaotic polynomial maps

XU ZHANG [1]

Department of Mathematics, Michigan State University, East Lansing, MI 48824, USA

Abstract. This paper introduces a class of polynomial maps in Euclidean spaces, investigates the conditions under which there exist Smale horseshoes and uniformly hyperbolic invariant sets, studies the chaotic dynamical behavior and strange attractors, and shows that some maps are chaotic in the sense of Li-Yorke or Devaney. This type of maps includes both the Logistic map and the Hénon map. For some maps in three-dimensional spaces under certain conditions, if the expansion dimension is equal to one or two, it is shown that there exist a Smale horseshoe and a uniformly hyperbolic invariant set on which the system is topologically conjugate to the two-sided fullshift on finite alphabet; if the system is expanding, then it is verified that there is an forward invariant set on which the system is topologically semi-conjugate to the one-sided fullshift on eight symbols. For three types of high-dimensional polynomial maps with degree two, the existence of Smale horseshoe and the uniformly hyperbolic invariant sets are studied, and it is proved that the map is topologically conjugate to the two-sided fullshift on finite alphabet on the invariant set under certain conditions. Some interesting maps with chaotic attractors and positive Lyapunov exponents in three-dimensional spaces are found by using computer simulations. In the end, two examples are provided to illustrate the theoretical results.

Keywords: Attractor; Devaney chaos; fullshift; Hénon map; Li-Yorke chaos; Logistic map; polynomial map; Smale horseshoe; uniformly hyperbolic set.

1 Introduction

The development of chaos theory dates back to Poincaré's work on the three-body problem [40, 41]. Thanks to the discovery of the Lorenz attractor and Li-Yorke chaos [33, 34, 45], the chaos theory is gradually becoming popular. Chaos theory is being successfully applied in many fields, from natural science to engineering. For example, it is an important branch of

---

[1] Email address: xuzhang08@gmail.com (X. Zhang).

dynamical systems in mathematics [51], planetary orbits [38] and fluid motion [19] in physics, the analysis of chemical compounds [48] in chemistry, secure communication [21] and chaos control [13] in engineering, new types of econometric models [30] and job selection [42] in economics.

The well-known chaotic polynomial maps are the Logistic map and the Hénon map, which were brought forward by May [35] and Hénon [28], respectively. These examples are important models in the research of nonlinear dynamics because of their simple expressions and complicated dynamical behavior. In engineering, chaotic polynomial maps were used as random number generators and have many applications [49], such as spread spectrum communications [31], and cryptosystems containing image encoding [8] and data encryption [29]. Since the encoding methods are dependent on the properties of the chaotic maps, and the expressions for the polynomial maps are relatively easy, the polynomial maps in our present work might be very useful in the future applications.

The Hénon map or generalized Hénon map is an important model in the research of two-dimensional polynomial diffeomorphic maps. Devaney and Nitecki investigated the conditions under which the real quadratic Hénon map has a hyperbolic invariant set on which it is topologically conjugate to the two-sided fullshift on two symbols [17]. Friedland and Milnor showed that the polynomial diffeomorphic map from the real or complex plane to itself is either conjugate to a composition of generalized Hénon maps or dynamically trivial [23]. Dullin and Meiss studied the conditions which imply that the real cubic Hénon map has hyperbolic invariant set [18]. The deeply relationship between the dynamical behavior and the change of parameters for the real Hénon map was described clearly based on the work contributed by Benedicks, Carleson, Viana, Young and et al. [9, 10, 11]. Bedford and Smillie applied the techniques in complex dynamics to show the existence of a hyperbolic horseshoe and a quadratic tangency between stable and unstable manifolds of fixed points for the real Hénon map under certain conditions, where the parameter is on the boundary of the set, in which the map is uniformly hyperbolic on its non-wandering set [6, 7]. Cao et al. investigated the existence of the uniformly hyperbolic set and an orbit of tangency for the Hénon-like families of diffeomorphisms of real plane by using the methods in real dynamics [12]. The hyperbolicity, ergodicity, Lyapunov exponents, topological entropy and so on, were also discussed for the complex Hénon maps (see [3, 4, 5] and references therein). In [52], we studied the existence of Smale horseshoe and uniformly hyperbolic invariant set on which the map is topologically conjugate to the two-sided fullshift on finite alphabet for the generalized Hénon maps.

There are many interesting chaotic attractors have been found in the study of the three-dimensional polynomial maps. Many authors investigated the following three-dimensional map $F : (x, y, z) \in \mathbb{R}^3 \to (f_1, f_2, f_3) \in \mathbb{R}^3$:

$$\begin{cases} f_1(x,y,z) = y \\ f_2(x,y,z) = z \\ f_3(x,y,z) = M_1 + Bx + M_2 y - z^2 \end{cases} \tag{1.1}$$

where M_1, M_2, B are parameters. In [26], Gonchenko et.al. studied the wild Lorenz-type strange attractors of system (1.1). In [25], Gonchenko et.al. studied the existence of wild-hyperbolic strange attractors and homoclinic bifurcations of system (1.1). In [27], Gonchenko et.al. investigated the bifurcations of system (1.1) with non-transverse heteroclinic cycles. In [32], Li and Yang investigated the three-dimensional Smale horseshoe and gave a computer assisted verification of the existence of hyperchaos of a class of three-dimensional polynomial maps. In [20], Elhadj and Sprott determined all the possible forms of the three-dimensional quadratic diffeomorphisms with constant Jacobian. In [22], Fournier-Prunaret et.al. studied the bifurcation and chaotic dynamics of a kind of three-dimensional maps of logistic type. In [24], Gonchenko and Li studied the existence of the uniformly hyperbolic invariant sets topologically conjugating to the Smale horseshoe for two cases of three-dimensional quadratic maps with constant Jacobian. In [46, 47], Sprott gave the idea that the computer simulations play an important role in the search of the chaotic attractors and studied several types of polynomial maps.

The polynomial diffeomorphic maps in high-dimensional spaces are different from those in two-dimensional spaces, since the degree of the inverse maps might be a large number and it is difficult to find the explicit expressions for the inverse maps. For more examples, please refer to the maps in Section 3 or related references [37]. There exists a bound on the degree of the inverse, that is, degree$(F^{-1}) \leq (\text{degree}(F))^{n-1}$ [14]. To investigate the inverse of the polynomial maps, an interesting problem is the well-known Jacobian Conjecture on $\mathbb{R}^n$, that is, is every polynomial map $F : \mathbb{R}^n \to \mathbb{R}^n$ with non-zero constant det$F'(x)$ a bijective map with a polynomial inverse? This is called the real Jacobian Conjecture, if the map is defined on $\mathbb{C}^n$, then this is the complex Jacobian Conjecture. The important progress on this problem includes Moh's proof for $n = 2$ and degree$(F) \leq 100$ [36], Wang's study for all n and degree$(F) \leq 2$ [50], Bass, Connell, and Wright reduced the problem to a special case, and obtained that the maps of homogeneous type of degree exactly three can imply the Jacobian Conjecture [2], and so on.

We introduce the following type of maps:

$$\begin{cases} f_1(x_1, ..., x_n) = a_{11}p_1(x_1) + \sum_{2\leq j\leq n} a_{1j}x_j \\ \vdots \qquad\quad \vdots \qquad\quad \vdots \qquad\quad \vdots \\ f_k(x_1, ..., x_n) = a_{kk}p_k(x_k) + \sum_{1\leq j\leq n,\ j\neq k} a_{kj}x_j \\ \vdots \qquad\quad \vdots \qquad\quad \vdots \qquad\quad \vdots \\ f_n(x_1, ..., x_n) = a_{nn}p_n(x_n) + \sum_{1\leq j\leq n-1} a_{nj}x_j \end{cases}, \tag{1.2}$$

where $p_i(x_i)$, $1 \leq i \leq n$, are real polynomials, a_{ij}, $1 \leq i, j \leq n$, are real parameters. We study the case that $n \geq 3$. For $n = 1, 2$, some results can be found in [52, 53].

First, we study the following type of three-dimensional polynomial maps $F : (x, y, z) \in \mathbb{R}^3 \to (f_1, f_2, f_3) \in \mathbb{R}^3$:

$$\begin{cases} f_1(x, y, z) = a_1p(x) + a_2y + a_3z \\ f_2(x, y, z) = b_1x + b_2q(y) + b_3z \\ f_3(x, y, z) = c_1x + c_2y + c_3r(z), \end{cases} \tag{1.3}$$

where a_i, b_i, c_i are real parameters, $1 \leq i \leq 3$, $p(x)$, $q(y)$, and $r(z)$ are real polynomials.

(1). We investigate a type of diffeomorphism with polynomial inverse by assuming that $a_1 \neq 0$, $b_2 = c_3 = 0$, and $p(x)$ having two different non-negative real zeros, and verify the existence of a Smale horseshoe and a uniformly hyperbolic invariant set on which the system is topologically conjugate to the two-sided fullshift on two symbols under certain conditions (See Theorems 3.1–3.2). If $p(x)$ has only m simple real roots, then there is a uniformly hyperbolic invariant set on which the system is topologically conjugate to the two-sided fullshift on m symbols under certain conditions (see Theorem 3.3).

(2). We study a class of diffeomorphism with polynomial inverse, where $a_1b_2 \neq 0$, $c_3 = 0$, $p(x)$ and $q(y)$ are assumed to have two distinct non-negative real roots, and show that there exist a Smale horseshoe and a uniformly hyperbolic invariant set on which the system is topologically conjugate to the two-sided fullshift on four symbols under certain conditions (See Theorems 3.4– 3.6). If $p(x)$ has only m simple real roots, and $q(y)$ has only n simple real zeros, then there is a uniformly hyperbolic invariant set on which the system is topologically conjugate to the two-sided fullshift on mn symbols under certain conditions (see Theorem 3.7).

(3). We study a kind of maps, where $a_1 b_2 c_3 \neq 0$, $p(x)$, $q(y)$, and $r(z)$ have two different non-negative real roots, and prove that there is an forward invariant set on which the system is topologically semi-conjugate to the one-sided fullshift on eight symbols (See Theorems 3.8–3.11).

Second, we investigate the following type of maps : $F : (x_1, ..., x_n) \in \mathbb{R}^n \to (f_1, ..., f_n) \in \mathbb{R}^n$:

$$\begin{cases} f_1(x_1, ..., x_n) = a_{11} - x_1^2 + \sum_{2 \leq j \leq n} a_{1j} x_j \\ \vdots \qquad\quad \vdots \qquad\quad \vdots \qquad\quad \vdots \\ f_d(x_1, ..., x_n) = a_{dd} - x_d^2 + \sum_{1 \leq j \leq n,\ j \neq d} a_{dj} x_j \\ f_{d+1}(x_1, ..., x_n) = \sum_{1 \leq j \leq n,\ j \neq d+1} a_{d+1,j} x_j \\ \vdots \qquad\quad \vdots \qquad\quad \vdots \qquad\quad \vdots \\ f_n(x_1, ..., x_n) = \sum_{1 \leq j \leq n-1} a_{nj} x_j \end{cases}, \tag{1.4}$$

where a_{ij}, $1 \leq i, j \leq n$, are real parameters. We study three kinds of polynomial maps by giving the inverse expressions, and show that there exist Smale horseshoe and uniformly hyperbolic invariant sets on which the maps are topologically conjugate with fullshift on finite alphabet (see Theorem 4.1).

We show that some maps are chaotic in the sense of Li-Yorke or Devaney. We also apply the computer simulation methods to study the existence of chaotic attractors and calculate the maximal Lyapunov exponents (see Section 5).

The rest of the paper is organized as follows. In Section 2, we introduce some basic concepts and results. In Section 3, we investigate the existence of uniformly hyperbolic invariant sets and complicated dynamics of maps (1.3) in three-dimensional spaces. We split this section into three parts. In Section 4, we study the existence of uniformly hyperbolic invariant sets and the chaotic dynamics of the maps (1.4) in high-dimensional Euclidean spaces. In Section 5, some interesting maps with chaotic attractors and positive Lyapunov exponents are obtained by applying computer simulations. In Section 6, several examples are given to illustrate the theoretical results.

2 Preliminaries

In this section, the concepts of uniformly hyperbolic set, symbolic dynamics, and chaos, and related results are introduced.

Definition 2.1. [39, 43] Consider a map $F : \mathbb{R}^n \to \mathbb{R}^n$. An invariant set $\Lambda \subset \mathbb{R}^n$ is called a uniformly hyperbolic set for F, if there are constants $C > 0$, $\lambda > 1$, and a continuous splitting $T_w\mathbb{R}^n = E^u_w \oplus E^s_w$, $w \in \Lambda$, such that

1 . $DF_w(E^s_w) = E^s_{F(w)}$ and $DF_w(E^u_w) = E^u_{F(w)}$;

2 . $|DF^m_w(v)| \geq C\lambda^m|v|$, $v \in E^u_w$, $m \geq 0$;

3 . $|DF^{-m}_w(v)| \geq C\lambda^m|v|$, $v \in E^s_w$, $m \geq 0$.

Given a subspace $E \subset \mathbb{R}^n$ with $0 < \dim E < n$ and $\mathbb{R}^n = E \oplus E^c$, where E^c is the algebraic complement of E, the standard unit cone with respect to E is given by:

$$K_1(E, E^c) = \left\{ \mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} : \ v_1 \in E, \ v_2 \in E^c, \text{ and } |v_2| \leq |v_1| \right\}.$$

Given a linear automorphism $T : \mathbb{R}^n \to \mathbb{R}^n$ with $T(E) = E$, the image $T(K_1(E, E^c))$ is called a cone in $\mathbb{R}^n$ with core E, denoted by $\mathcal{C}(E)$. For convenience, $\mathcal{C}$ is used to represent a cone in $\mathbb{R}^n$, where the proper subspace E of $\mathbb{R}^n$ is omitted.

Suppose that F is a diffeomorphism on $\mathbb{R}^n$ and U is a compact subset of $\mathbb{R}^n$, then $\Lambda = \cap^{\infty}_{-\infty} F^i(U)$ is a compact invariant set. Let the cone field $\mathcal{C} = \{\mathcal{C}_w\}$ on Λ be the set of cones $\mathcal{C}_w \subset T\mathbb{R}^n = \mathbb{R}^n$, $w \in \Lambda$. It is said that the cone field $\mathcal{C}_w$ has constant orbit core dimension on Λ if $\dim E_w = \dim E_{F(w)}$, $w \in \Lambda$, where E_w and $E_{F(w)}$ are the cores of $\mathcal{C}_w$ and $\mathcal{C}_{F(w)}$, respectively.

Set

$$m_{\mathcal{C},w} = m_{\mathcal{C},w}(F) := \inf_{\mathbf{v} \in \mathcal{C}_w \setminus \{0\}} \frac{|DF_w(\mathbf{v})|}{|\mathbf{v}|}$$

and

$$m'_{\mathcal{C},w} = m'_{\mathcal{C},w}(F) := \inf_{\mathbf{v} \notin \mathcal{C}_{F(w)}} \frac{|DF^{-1}_{F(w)}(\mathbf{v})|}{|\mathbf{v}|},$$

where $m_{\mathcal{C},w}$ is said to be the minimal expansion of F on $\mathcal{C}_w$, and $m'_{\mathcal{C},w}$ is called the minimal co-expansion of F on $\mathcal{C}_w$.

Lemma 2.1. ([39], Theorem 1.4) A necessary and sufficient condition for Λ to be a uniformly hyperbolic set for F is that there are an integer $N > 0$ and a cone field $\mathcal{C}$ with constant orbit core dimension over Λ such that F^N is both expanding and co-expanding on $\mathcal{C}$.

Next, we introduce the symbolic dynamics [43]. Set $S_0 := \{1, 2, ..., m\},\ m \geq 2$. Let

$$\sum\nolimits_m := \{\alpha = (..., a_{-2}, a_{-1}, a_0, a_1, a_2, ...) : \ a_i \in S_0, \quad i \in \mathbb{Z}\}$$

be the two-sided sequence space, where the distance on $\sum_m$ is defined by $d(\alpha, \beta) = \sum_{i=-\infty}^{\infty} \frac{d(a_i, b_i)}{2^{|i|}}$, for $\alpha = (..., a_{-2}, a_{-1}, a_0, a_1, a_2, ...)$, $\beta = (..., b_{-2}, b_{-1}, b_0, b_1, b_2, ...) \in \sum_m$, and $d(a_i, b_i) = 1$ if $a_i \neq b_i$, and $d(a_i, b_i) = 0$ if $a_i = b_i$, $i \in \mathbb{Z}$. The shift map $\sigma : \sum_m \to \sum_m$ is defined by $\sigma(\alpha) = (..., b_{-2}, b_{-1}, b_0, b_1, b_2...)$ for any $\alpha = (..., a_{-2}, a_{-1}, a_0, a_1, a_2...) \in \sum_m$, where $b_i = a_{i+1}$ for any $i \in \mathbb{Z}$. We call $(\sum_m, \sigma)$ is the two-sided symbolic dynamical system on m symbols, or simply two-sided fullshift on m symbols.

Two definitions of chaos are introduced, which will be useful in the sequel.

Definition 2.2. [33] Let (X, d) be a metric space, $F : X \to X$ a map, and S a subset of X with at least two points. Then S is called a scrambled set of F if for any two distinct points $x, y \in S$,

$$\liminf_{n\to\infty} d(F^n(x), F^n(y)) = 0, \ \limsup_{n\to\infty} d(F^n(x), F^n(y)) > 0.$$

The map F is said to be chaotic in the sense of Li-Yorke if there exists an uncountable scrambled set S of F.

Let (X, d) be a metric space. A continuous map $F : X \to X$ is said to be topologically transitive if for any two open subsets U and V of X, there exists a positive integer m such that $F^m(U) \cap V \neq \emptyset$; F is said to have sensitive dependence on initial conditions in X if there exists a positive constant δ such that for any point $x \in X$ and any neighborhood U of x, there exist $y \in U$ and a positive integer m such that $d(F^m(x), F^m(y)) > \delta$.

Definition 2.3. [16] Let (X, d) be a metric space. A map $F : V \subset X \to V$ is said to be chaotic on V in the sense of Devaney if

(i) the set of the periodic points of F is dense in V;

(ii) F is topologically transitive in V;

(iii) F has sensitive dependence on initial conditions in V.

In the above definition, condition (iii) is redundant if F is continuous in V by the result of [1], where V is an infinite set.

Lemma 2.2. [44, Theorem 2.2] The fullshift on finite alphabet is chaotic in the sense of both of Li-Yorke and Devaney.

3 Smale horseshoe in three-dimensional space

In this section, we study the parameter regions of the maps (1.3) such that there exist invariant sets on which the maps are uniformly hyperbolic and topologically conjugate with fullshift on finite alphabet.

For map (1.3), consider the situations that the polynomials $p(x)$, $q(y)$, and $r(z)$ have at least two distinct real zeros.

The polynomial p can be written as

$$p(x) = (x-\alpha_1)^{m_1}\cdots(x-\alpha_{r_1})^{m_{r_1}}(y^2+\beta_{r_1+1}y+\gamma_{r_1+1})^{m_{r_1+1}}\cdots(y^2+\beta_{s_1}y+\gamma_{s_1})^{m_{s_1}},$$

where

$$m_i \geq 1, 1 \leq i \leq s_1;\ \sum_{i=1}^{r_1} m_i + \sum_{i=r_1+1}^{s_1} 2m_i = d_1;\ \beta_j^2 - 4\gamma_j < 0,\ r_1+1 \leq j \leq s_1,$$

and $\alpha_1, ..., \alpha_{r_1}$ are real zeros of $p(x)$ with $\alpha_1 < \cdots < \alpha_{r_1}$.

The polynomial $q(y)$ can be represented as follows:

$$q(y) = (y-\xi_1)^{n_1}\cdots(y-\xi_{r_2})^{n_{r_2}}(y^2+\eta_{r_2+1}y+\zeta_{r_2+1})^{n_{r_2+1}}\cdots(y^2+\eta_{s_2}y+\zeta_{s_2})^{n_{s_2}},$$

where

$$n_i \geq 1, 1 \leq i \leq s_2;\ \sum_{i=1}^{r_2} n_i + \sum_{i=r_2+1}^{s_2} 2n_i = d_2;\ \eta_j^2 - 4\zeta_j < 0,\ r_2+1 \leq j \leq s_2,$$

and $\xi_1, ..., \xi_{r_2}$ are real zeros of $q(y)$ with $\xi_1 < \cdots < \xi_{r_2}$.

The polynomial $r(z)$ can be expressed as follows:

$$r(z) = (z-\kappa_1)^{l_1}\cdots(z-\kappa_{r_3})^{l_{r_3}}(z^2+\mu_{r_3+1}z+\nu_{r_3+1})^{l_{r_3+1}}\cdots(z^2+\mu_{s_3}z+\nu_{s_3})^{l_{s_3}},$$

where

$$l_i \geq 1, 1 \leq i \leq s_3;\ \sum_{i=1}^{r_3} l_i + \sum_{i=r_3+1}^{s_3} 2l_i = d_3;\ \mu_k^2 - 4\nu_k < 0,\ r_3+1 \leq k \leq s_3,$$

and $\kappa_1, ..., \kappa_{r_3}$ are real zeros of $r(z)$ with $\kappa_1 < \cdots < \kappa_{r_3}$.

It is easy to obtain that

$$\frac{p'(x)}{p(x)} = \sum_{i=1}^{r_1} \frac{m_i}{x-\alpha_i} + \sum_{i=r_1+1}^{s_1} \frac{m_i(2x+\beta_i)}{x^2+\beta_i x+\gamma_i},$$

$$\frac{q'(y)}{q(y)} = \sum_{i=1}^{r_2} \frac{n_i}{y-\xi_i} + \sum_{i=r_2+1}^{s_2} \frac{n_i(2y+\eta_i)}{y^2+\eta_i y+\zeta_i}.$$

$$\frac{r'(z)}{r(z)} = \sum_{i=1}^{r_3} \frac{l_i}{z-\kappa_i} + \sum_{i=r_3+1}^{s_3} \frac{l_i(2z+\mu_i)}{z^2+\mu_i z+\nu_i}.$$

So,

$$\lim_{x\to\alpha_{i_0}^+} \frac{p'(x)}{p(x)} = \infty, \quad \lim_{x\to\alpha_{i_0+1}^-} \frac{p'(x)}{p(x)} = -\infty; \tag{3.1}$$

$$\lim_{y\to\xi_{j_0}^+} \frac{q'(y)}{q(y)} = \infty, \quad \lim_{y\to\xi_{j_0+1}^-} \frac{q'(y)}{q(y)} = -\infty; \tag{3.2}$$

$$\lim_{z\to\kappa_{k_0}^+} \frac{r'(z)}{r(z)} = \infty, \quad \lim_{z\to\kappa_{k_0+1}^-} \frac{r'(z)}{r(z)} = -\infty. \tag{3.3}$$

Lemma 3.1. [52, Lemma 3.1]

(i) Suppose that there exists i_0, $1 \le i_0 < r_1$, such that $\alpha_{i_0} \ge 0$. Then, there exist $\delta_1, \delta_1' \in (\alpha_{i_0}, \alpha_{i_0+1})$ such that $p^{(m_{i_0})}(\alpha_{i_0})p'(x) > 0$ for all $x \in (\alpha_{i_0}, \delta_1]$, $p^{(m_{i_0})}(\alpha_{i_0})p'(x) < 0$ for all $x \in [\delta_1', \alpha_{i_0+1})$.

(ii) Suppose that there exists j_0, $1 \le j_0 < r_2$, such that $\xi_{j_0} \ge 0$. Then, there exist $\delta_2, \delta_2' \in (\xi_{j_0}, \xi_{j_0+1})$ such that $q^{(n_{j_0})}(\xi_{j_0})q'(y) > 0$ for all $y \in (\xi_{j_0}, \delta_2]$, $q^{(n_{j_0})}(\xi_{j_0})q'(y) < 0$ for all $y \in [\delta_2', \xi_{j_0+1})$.

(iii) Suppose that there exists k_0, $1 \le k_0 < r_3$, such that $\kappa_{k_0} \ge 0$. Then, there exist $\delta_3, \delta_3' \in (\kappa_{k_0}, \kappa_{k_0+1})$ such that $r^{(l_{k_0})}(\kappa_{k_0})r'(z) > 0$ for all $z \in (\kappa_{k_0}, \delta_3]$, $r^{(l_{k_0})}(\kappa_{k_0})r'(z) < 0$ for all $z \in [\delta_3', \kappa_{k_0+1})$.

3.1 The maps with the dimension of the unstable subspace equals to one

In this subsection, we investigate the polynomial maps (1.3) in the following form, $F : (x, y, z) \in \mathbb{R}^3 \to (f_1, f_2, f_3) \in \mathbb{R}^3$:

$$\left\{ \begin{array}{l} f_1(x,y,z) = a_1 p(x) + a_2 y + a_3 z \\ f_2(x,y,z) = b_1 x \\ f_3(x,y,z) = c_2 y \end{array} \right. , \tag{3.4}$$

where a_1, a_2, a_3, b_1, c_2 are real parameters and $a_3 b_1 c_2 \neq 0$.

The derivative of (3.4) is

$$DF = \begin{pmatrix} a_1 p'(x) & a_2 & a_3 \\ b_1 & 0 & 0 \\ 0 & c_2 & 0 \end{pmatrix}. \tag{3.5}$$

The inverse of F is $H : (x, y, z) \in \mathbb{R}^3 \to (h_1, h_2, h_3) \in \mathbb{R}^3$:

$$\left\{ \begin{array}{l} h_1(x,y,z) = \frac{y}{b_1} \\ h_2(x,y,z) = \frac{z}{c_2} \\ h_3(x,y,z) = \frac{1}{a_3}\Big(x - a_1 p(y/b_1) - \frac{a_2}{c_2} z \Big) \end{array} \right. \tag{3.6}$$

and the derivative of H is

$$DF^{-1} = \begin{pmatrix} 0 & \frac{1}{b_1} & 0 \\ 0 & 0 & \frac{1}{c_2} \\ \frac{1}{a_3} & -\frac{a_1}{b_1 a_3} p'(y/b_1) & -\frac{a_2}{a_3 c_2} \end{pmatrix}. \tag{3.7}$$

So, the determinant of the Jacobian of this type of maps is constant.

To start our work, an example is introduced as follows:

$$\left\{ \begin{array}{l} f_1(x,y,z) = a_1 x(1-x) + a_2 y + a_3 z \\ f_2(x,y,z) = b_1 x \\ f_3(x,y,z) = c_2 y \end{array} \right. .$$

This example can be thought of as the generalization of the Hénon map in three-dimensional spaces.

Theorem 3.1. Suppose that there are $1 \leq i_0 < r_1$ such that $0 = \alpha_{i_0} < \alpha_{i_0+1}$ and $m_{i_0} = m_{i_0+1} = 1$, that is, there are two distinct non-negative zeros of $p(x)$ with the multiplicity equals to one. If

$$\max\{a_3 c_2 b_1 \alpha_{i_0+1}, a_2 b_1 \alpha_{i_0+1}, a_2 b_1 \alpha_{i_0+1} + a_3 c_2 b_1 \alpha_{i_0+1}\} \leq 0 \text{ and } a_1 p'(0) > 0, \tag{3.8}$$

then, for fixed a_2, a_3, b_1, and c_2 with $|c_2| < 1$, and sufficiently large $|a_1|$, there exist a Smale horseshoe and a uniformly hyperbolic invariant set Λ for the map (3.4), on which F is topologically conjugate to two-sided fullshift on two symbols. Consequently, they are chaotic in the sense of both Li-Yorke and Devaney.

For convenient, assume that $[a, b]$ and $[b, a]$ represent the same interval. Set

$$U := [0, \alpha_{i_0+1}] \times [0, b_1\alpha_{i_0+1}] \times [0, c_2b_1\alpha_{i_0+1}],\ \ \lambda := \frac{1}{|c_2|}.$$

Lemma 3.2. Under the assumptions of Theorem 3.1, $F(U) \cap U$ and $F^{-1}(U) \cap U$ have two non-empty connected components, respectively.

Proof. First, we need to determine the image of the vertices of the cubic U under F. The vertices of U are

$$A = (\alpha_{i_0+1}, 0, 0), B = (\alpha_{i_0+1}, b_1\alpha_{i_0+1}, 0), C = (0, b_1\alpha_{i_0+1}, 0), D = (0, 0, 0),$$

$$A' = (\alpha_{i_0+1}, 0, c_2b_1\alpha_{i_0+1}), B' = (\alpha_{i_0+1}, b_1\alpha_{i_0+1}, c_2b_1\alpha_{i_0+1}),$$

$$C' = (0, b_1\alpha_{i_0+1}, c_2b_1\alpha_{i_0+1}), D' = (0, 0, c_2b_1\alpha_{i_0+1}).$$

The images of these vertices under F are

$$F(A) = (0, b_1\alpha_{i_0+1}, 0), F(B) = (a_2b_1\alpha_{i_0+1}, b_1\alpha_{i_0+1}, c_2b_1\alpha_{i_0+1}),$$

$$F(C) = (a_2b_1\alpha_{i_0+1}, 0, c_2b_1\alpha_{i_0+1}), F(D) = (0, 0, 0),$$

$$F(A') = (a_3c_2b_1\alpha_{i_0+1}, b_1\alpha_{i_0+1}, 0), F(B') = (a_2b_1\alpha_{i_0+1} + a_3c_2b_1\alpha_{i_0+1}, b_1\alpha_{i_0+1}, c_2b_1\alpha_{i_0+1}),$$

$$F(C') = (a_2b_1\alpha_{i_0+1} + a_3c_2b_1\alpha_{i_0+1}, 0, c_2b_1\alpha_{i_0+1}), F(D') = (a_3c_2b_1\alpha_{i_0+1}, 0, 0).$$

From the map (3.4), it follows that the transformation is linear with respect to the y and z variables. For the x variable, it is a parabolic-like function for $x \in [0, \alpha_{i_0+1}]$. By (3.8) and the calculation of the vertices above, one has that if $|a_1|$ is sufficiently large, then there are two connected components of $U \cap F(U)$.

On the other hand, the image of the vertices under $F^{-1} = H$ are

$$H(A) = (0, 0, \alpha_{i_0+1}/a_3), H(B) = (\alpha_{i_0+1}, 0, \alpha_{i_0+1}/a_3), H(C) = (\alpha_{i_0+1}, 0, 0), H(D) = (0, 0, 0),$$

$$H(A') = (0, b_1\alpha_{i_0+1}, \alpha_{i_0+1}(1 - a_2b_1)/a_3), H(B') = (\alpha_{i_0+1}, b_1\alpha_{i_0+1}, \alpha_{i_0+1}(1 - a_2b_1)/a_3),$$

$$H(C')=(\alpha_{i_0+1},b_1\alpha_{i_0+1},-a_2b_1\alpha_{i_0+1}/a_3),H(D')=(0,b_1\alpha_{i_0+1},-a_2b_1\alpha_{i_0+1}/a_3).$$

Since F is diffeomorphic, $U\cap F^{-1}(U)=F^{-1}(U\cap F(U))$ has two connected components. This completes the proof. □

Lemma 3.3. Under the assumptions of Theorem 3.1, the invariant set Λ is uniformly hyperbolic.

Proof. By (i) of Lemma 3.1, assume that $|a_1|$ is large enough such that for $w=(x,y,z)\in U\cap F^{-1}(U)$, one has that $x\in[0,\delta_1]\cup[\delta_1',\alpha_{i_0+1}]$.

Since $m_{i_0}=m_{i_0+1}=1$, set

$$M_0:=\min_{x\in[0,\delta_1]\cup[\delta_1',\alpha_{i_0+1}]}|p'(x)|>0.$$

Suppose that

$$|a_1|\geq\max\left\{\frac{\lambda+|a_2|+|a_3|}{M_0},\ \frac{\lambda|a_3b_1c_2|+|b_1c_2|+|a_2b_1|}{M_0|c_2|}\right\}. \tag{3.9}$$

Now, it is to show that for any $w_0\in U\cap F^{-1}(U)$, one has that $m_{\mathcal{C},w_0}\geq\lambda$.

We introduce the unit cone:

$$K_1(\mathbb{R},\mathbb{R}^2)=\left\{\mathbf{v}=\begin{pmatrix}v_0\\u_0\end{pmatrix}:\ v_0\in\mathbb{R},\ u_0=(u_{01},u_{02})^T\in\mathbb{R}^2,\ \text{and }|u_0|\leq|v_0|\right\},$$

where $|u_0|=\max\{|u_{01}|,|u_{02}|\}$ and $|\mathbf{v}|=\max\{|v_0|,|u_0|\}$.

Denote $\begin{pmatrix}v_1\\u_1\end{pmatrix}=DF_{w_0}\begin{pmatrix}v_0\\u_0\end{pmatrix}$, where $\begin{pmatrix}v_0\\u_0\end{pmatrix}\in K_1(\mathbb{R},\mathbb{R}^2)$. By (3.5),

$$\begin{cases}v_1=a_1p'(x_0)v_0+a_2u_{01}+a_3u_{02}\\u_{11}=b_1v_0\\u_{12}=c_2u_{01}\end{cases},$$

this, together with (3.9), implies that

$$|v_1|\geq|v_0||a_1p'(x_0)|-|a_2||u_{01}|-|a_3||u_{02}|\geq|v_0||a_1p'(x_0)|-|a_2||v_0|-|a_3||v_0|\geq\lambda|v_0|.$$

Hence, $m_{\mathcal{C},w_0}\geq\lambda>1$.

Next, it is to show that for any point $w_0\in U\cap F(U)$, $m'_{\mathcal{C},w_0}\geq\lambda$. For any $\begin{pmatrix}v_0\\u_0\end{pmatrix}\notin K_1(\mathbb{R},\mathbb{R}^2)$, denote $\begin{pmatrix}v_{-1}\\u_{-1}\end{pmatrix}=DF^{-1}_{w_0}\begin{pmatrix}v_0\\u_0\end{pmatrix}$, it follows from (3.7) that

$$\begin{cases}v_{-1}=\frac{1}{b_1}u_{01}\\u_{-11}=\frac{1}{c_2}u_{02}\\u_{-12}=\frac{1}{a_3}v_0-\frac{a_1}{b_1a_3}p'(y_0/b_1)u_{01}-\frac{a_2}{a_3c_2}u_{02}\end{cases}.$$

If $|u_0| = |u_{01}|$, then by (3.9),

$$\begin{aligned}|u_{-1}| \geq& |u_{-12}| \geq -\frac{1}{|a_3|}|v_0| + \frac{1}{|b_1a_3|}|a_1p'(y_0/b_1)||u_{01}| - \frac{|a_2|}{|a_3c_2|}|u_{02}|\\ \geq& \frac{1}{|b_1a_3|}|a_1p'(y_0/b_1)||u_0| - \frac{1}{|a_3|}|u_0| - \frac{|a_2|}{|a_3c_2|}|u_0|\\ \geq& \lambda|u_0|;\end{aligned}$$

if $|u_0| = |u_{02}|$, then

$$|u_{-1}| \geq |u_{02}/c_2| = \lambda|u_{0,2}| = \lambda|u_0|.$$

Hence, $m'_{\mathcal{C},w_0} \geq \lambda$.

Thus, by Lemma 2.1, one has that the invariant set Λ is uniformly hyperbolic. The proof is completed. □

Therefore, it follows from Lemmas 2.2, 3.2, and 3.3 that Theorem 3.1 holds.

Theorem 3.2. Suppose that there are $1 \leq i_0 < r_1$ such that $0 < \alpha_{i_0} < \alpha_{i_0+1}$, that is, there are two distinct positive zeros of $p(x)$. Set

$$M_1 := \max\{a_2b_1\alpha_{i_0}+a_3c_2b_1\alpha_{i_0}, a_2b_1\alpha_{i_0}+a_3c_2b_1\alpha_{i_0+1}, a_2b_1\alpha_{i_0+1}+a_3c_2b_1\alpha_{i_0}, a_2b_1\alpha_{i_0+1}+a_3c_2b_1\alpha_{i_0+1}\},$$

$$M_2 := \min\{a_2b_1\alpha_{i_0}+a_3c_2b_1\alpha_{i_0}, a_2b_1\alpha_{i_0}+a_3c_2b_1\alpha_{i_0+1}, a_2b_1\alpha_{i_0+1}+a_3c_2b_1\alpha_{i_0}, a_2b_1\alpha_{i_0+1}+a_3c_2b_1\alpha_{i_0+1}\}.$$

Consider the following two situations:

$$\text{(i). } M_1 < \alpha_{i_0} \text{ and } a_1p^{(m_{i_0})}(\alpha_{i_0}) > 0; \text{ (ii). } M_2 > \alpha_{i_0+1} \text{ and } a_1p^{(m_{i_0})}(\alpha_{i_0}) < 0. \tag{3.10}$$

Then, for fixed a_2, a_3, b_1, and c_2 with $|c_2| < 1$, and sufficiently large $|a_1|$, there exist a Smale horseshoe and a uniformly hyperbolic invariant set Λ for the map (3.4), on which F is topologically conjugate to two-sided fullshift on two symbols. Consequently, they are chaotic in the sense of both Li-Yorke and Devaney.

For convenient, suppose that $[a, b]$ and $[b, a]$ represent the same interval. Denote

$$U := [\alpha_{i_0}, \alpha_{i_0+1}] \times [b_1\alpha_{i_0}, b_1\alpha_{i_0+1}] \times [c_2b_1\alpha_{i_0}, c_2b_1\alpha_{i_0+1}].$$

Set $\lambda := \frac{1}{|c_2|}$. It follows from the assumption $0 < |c_2| < 1$ that $\lambda > 1$. The invariant set is given by

$$\Lambda := \bigcap_{-\infty}^{+\infty} F^i(U).$$

Lemma 3.4. In Cases (i)-(ii) of Theorem 3.2, for fixed a_2, a_3, b_1, and c_2, and sufficiently large $|a_1|$, one has that $F(U)\cap U$ and $F^{-1}(U)\cap U$ have two non-empty connected components, respectively.

Proof. By the definition of map (3.4), one has that the transformation is linear with respect to the y and z variables.

Next, it is to study the transformation with respect to the x variable. First, it is to obtain the image of the vertices of U under the map F. The vertices of U are

$$A = (\alpha_{i_0+1}, b_1\alpha_{i_0}, c_2b_1\alpha_{i_0}), B = (\alpha_{i_0+1}, b_1\alpha_{i_0+1}, c_2b_1\alpha_{i_0}),$$

$$C = (\alpha_{i_0}, b_1\alpha_{i_0+1}, c_2b_1\alpha_{i_0}), D = (\alpha_{i_0}, b_1\alpha_{i_0}, c_2b_1\alpha_{i_0}),$$

$$A' = (\alpha_{i_0+1}, b_1\alpha_{i_0}, c_2b_1\alpha_{i_0+1}), B' = (\alpha_{i_0+1}, b_1\alpha_{i_0+1}, c_2b_1\alpha_{i_0+1}),$$

$$C' = (\alpha_{i_0}, b_1\alpha_{i_0+1}, c_2b_1\alpha_{i_0+1}), D' = (\alpha_{i_0}, b_1\alpha_{i_0}, c_2b_1\alpha_{i_0+1}).$$

The images under F are

$$F(A) = (a_2b_1\alpha_{i_0}+a_3c_2b_1\alpha_{i_0}, b_1\alpha_{i_0+1}, c_2b_1\alpha_{i_0}), F(B) = (a_2b_1\alpha_{i_0+1}+a_3c_2b_1\alpha_{i_0}, b_1\alpha_{i_0+1}, c_2b_1\alpha_{i_0+1}),$$

$$F(C) = (a_2b_1\alpha_{i_0+1} + a_3c_2b_1\alpha_{i_0}, b_1\alpha_{i_0}, c_2b_1\alpha_{i_0+1}), F(D) = (a_2b_1\alpha_{i_0} + a_3c_2b_1\alpha_{i_0}, b_1\alpha_{i_0}, c_2b_1\alpha_{i_0}),$$

$$F(A') = (a_2b_1\alpha_{i_0}+a_3c_2b_1\alpha_{i_0+1}, b_1\alpha_{i_0+1}, c_2b_1\alpha_{i_0}), F(B') = (a_2b_1\alpha_{i_0+1}+a_3c_2b_1\alpha_{i_0+1}, b_1\alpha_{i_0+1}, c_2b_1\alpha_{i_0+1}),$$

$$F(C') = (a_2b_1\alpha_{i_0+1}+a_3c_2b_1\alpha_{i_0+1}, b_1\alpha_{i_0}, c_2b_1\alpha_{i_0+1}), F(D') = (a_2b_1\alpha_{i_0}+a_3c_2b_1\alpha_{i_0+1}, b_1\alpha_{i_0}, c_2b_1\alpha_{i_0}).$$

This, together with (3.10), implies that there are two connected components of $U \cap F(U)$ for sufficiently large $|a_1|$.

On the other hand, the image of the vertices of U under $F^{-1} = H$ are

$$H(A) = \left(\alpha_{i_0}, b_1\alpha_{i_0}, \frac{1}{a_3}(\alpha_{i_0+1} - a_2b_1\alpha_{i_0})\right), H(B) = \left(\alpha_{i_0+1}, b_1\alpha_{i_0}, \frac{1}{a_3}(\alpha_{i_0+1} - a_2b_1\alpha_{i_0})\right),$$

$$H(C) = \left(\alpha_{i_0+1}, b_1\alpha_{i_0}, \frac{1}{a_3}(\alpha_{i_0} - a_2b_1\alpha_{i_0})\right), H(D) = \left(\alpha_{i_0}, b_1\alpha_{i_0}, \frac{1}{a_3}(\alpha_{i_0} - a_2b_1\alpha_{i_0})\right),$$

$$H(A') = \left(\alpha_{i_0}, b_1\alpha_{i_0+1}, \frac{1}{a_3}(\alpha_{i_0+1}-a_2b_1\alpha_{i_0+1})\right), H(B') = \left(\alpha_{i_0+1}, b_1\alpha_{i_0+1}, \frac{1}{a_3}(\alpha_{i_0+1}-a_2b_1\alpha_{i_0+1})\right),$$

$$H(C') = \left(\alpha_{i_0+1}, b_1\alpha_{i_0+1}, \frac{1}{a_3}(\alpha_{i_0} - a_2b_1\alpha_{i_0+1})\right), H(D') = \left(\alpha_{i_0}, b_1\alpha_{i_0+1}, \frac{1}{a_3}(\alpha_{i_0} - a_2b_1\alpha_{i_0+1})\right).$$

It follows from the fact F is diffeomorphic that $U \cap F^{-1}(U) = F^{-1}(U \cap F(U))$ has two connected components. The proof is completed. □

Lemma 3.5. In Cases (i)-(ii) of Theorem 3.2, for fixed a_2, a_3, b_1, and c_2, and sufficiently large $|a_1|$, one has that the invariant set Λ is uniformly hyperbolic.

Proof. In the following discussions, we always assume that a_2, a_3, b_1, and c_2 are fixed, and $|a_1|$ are sufficiently large. By (i) of Lemma 3.1, assume that $|a_1|$ is sufficiently large such that for $w = (x, y, z) \in U \cap F^{-1}(U)$, one has that $x \in (\alpha_{i_0}, \delta_1] \cup [\delta_1', \alpha_{i_0+1})$. For any point $w_0 = (x_0, y_0, z_0) \in U$, set $w_1 = F(w_0) = (x_1, y_1, z_1)$ and $w_{-1} = F^{-1}(w_0) = (x_{-1}, y_{-1}, z_{-1})$.

First, by (3.1) and Lemma 3.1, one has that for $w = (x, y, z) \in U \cap F^{-1}(U)$,

$$\text{Case (i)}: \quad \frac{|a_1 p'(x)|}{|a_1 p(x)|} \geq \frac{\max\left\{\lambda + |a_2| + |a_3|,\ \frac{\lambda|a_3 b_1 c_2| + |b_1 c_2| + |a_2 b_1|}{|c_2|}\right\}}{\alpha_{i_0} - M_1}; \tag{3.11}$$

$$\text{Case (ii)}: \quad \frac{|a_1 p'(x)|}{|a_1 p(x)|} \geq \frac{\max\left\{\lambda + |a_2| + |a_3|,\ \frac{\lambda|a_3 b_1 c_2| + |b_1 c_2| + |a_2 b_1|}{|c_2|}\right\}}{M_2 - \alpha_{i_0+1}}. \tag{3.12}$$

For any point $w_0 \in U \cap F^{-1}(U)$, one has

$$\begin{cases} x_1 = a_1 p(x_0) + a_2 y_0 + a_3 z_0 \\ y_1 = b_1 x_0 \\ z_1 = c_2 x_0 \end{cases}.$$

that is, $a_1 p(x_0) = x_1 - a_2 y_0 - a_3 z_0$, implying that

$$\text{Case (i)}: \quad a_1 p(x_0) \geq \alpha_{i_0} - M_1; \ \text{Case (ii)}: \quad -a_1 p(x_0) \geq M_2 - \alpha_{i_0+1}. \tag{3.13}$$

For any point $w_0 \in U \cap F(U)$, by (3.6), one has,

$$\begin{cases} x_{-1} = \frac{y_0}{b_1} \\ y_{-1} = \frac{z_0}{c_2} \\ z_{-1} = \frac{1}{a_3}\left(x_0 - a_1 p(y_0/b_1) - \frac{a_2}{c_2} z_0\right) \end{cases},$$

that is, $a_1 p(x_{-1}) = a_1 p(y_0/b_1) = x_0 - a_3 z_{-1} - \frac{a_2}{c_2} z_0$, yielding that

$$\text{Case (i)}: \quad a_1 p(x_{-1}) \geq \alpha_{i_0} - M_1; \ \text{Case (ii)}: \quad -a_1 p(x_{-1}) \geq M_2 - \alpha_{i_0+1}. \tag{3.14}$$

Hence, it follows from (3.11) and (3.13) that for any point $w_0 \in U \cap F^{-1}(U)$,

$$\text{Cases (i) and (ii)}: \quad |a_1 p'(x_0)| \geq \lambda + |a_2| + |a_3|. \tag{3.15}$$

Now, it is to show that for any $w_0 \in U \cap F(U)$, one has that $m_{\mathcal{C}, w_0} \geq \lambda$.

Consider the unit cone

$$K_1(\mathbb{R},\mathbb{R}^2)=\left\{\mathbf{v}=\begin{pmatrix} v_0\\ u_0\end{pmatrix}:\ v_0\in\mathbb{R},\ u_0=(u_{01},u_{02})^T\in\mathbb{R}^2,\ \text{and } |u_0|\le|v_0|\right\},$$

where $|u_0|=\max\{|u_{01}|,|u_{02}|\}$ and $|\mathbf{v}|=\max\{|v_0|,|u_0|\}$.

Suppose that $\begin{pmatrix} v_1\\ u_1\end{pmatrix}=DF_{w_0}\begin{pmatrix} v_0\\ u_0\end{pmatrix}$, where $\begin{pmatrix} v_0\\ u_0\end{pmatrix}\in K_1(\mathbb{R},\mathbb{R}^2)$. By (3.5),

$$\begin{cases} v_1=a_1p'(x_0)v_0+a_2u_{01}+a_3u_{02}\\ u_{11}=b_1v_0\\ u_{12}=c_2u_{01}\end{cases},$$

this, together with (3.15), implies that

$$|v_1|\ge|v_0||a_1p'(x_0)|-|a_2||u_{01}|-|a_3||u_{02}|\ge|v_0||a_1p'(x_0)|-|a_2||v_0|-|a_3||v_0|\ge\lambda|v_0|.$$

Hence, $m_{\mathcal{C},w_0}\ge\lambda$.

By (3.12) and (3.14), one has that for any point $w_0\in U\cap F(U)$,

$$|a_1p'(x_{-1})|=|a_1p'(y_0/b_1)|\ge\frac{\lambda|a_3b_1c_2|+|b_1c_2|+|a_2b_1|}{|c_2|},\tag{3.16}$$

Next, it is to show that for any point $w_0\in U\cap F(U)$, $m'_{\mathcal{C},w_0}\ge\lambda$. For any $\begin{pmatrix} v_0\\ u_0\end{pmatrix}\notin K_1(\mathbb{R},\mathbb{R}^2)$, that is , $|u_0|>|v_0|$, suppose $\begin{pmatrix} v_{-1}\\ u_{-1}\end{pmatrix}=DF^{-1}_{w_0}\begin{pmatrix} v_0\\ u_0\end{pmatrix}$, by (3.7), one has

$$\begin{cases} v_{-1}=\frac{1}{b_1}u_{01}\\ u_{-11}=\frac{1}{c_2}u_{02}\\ u_{-12}=\frac{1}{a_3}v_0-\frac{a_1}{b_1a_3}p'(y_0/b_1)u_{01}-\frac{a_2}{a_3c_2}u_{02}\end{cases}.$$

If $|u_0|=|u_{02}|$, then $|u_{-1}|\ge|u_{02}/c_2|=\lambda|u_{0,2}|=\lambda|u_0|$. If $|u_0|=|u_{01}|$, then by (3.16),

$$\begin{aligned}|u_{-1}|\ge&|u_{-12}|\ge-\frac{1}{|a_3|}|v_0|+\frac{1}{|b_1a_3|}|a_1p'(y_0/b_1)||u_{01}|-\frac{|a_2|}{|a_3c_2|}|u_{02}|\\ \ge&\frac{1}{|b_1a_3|}|a_1p'(y_0/b_1)||u_0|-\frac{1}{|a_3|}|u_0|-\frac{|a_2|}{|a_3c_2|}|u_0|\\ \ge&\lambda|u_0|.\end{aligned}$$

So, $m'_{\mathcal{C},w_0}\ge\lambda$.

Thus, it follows from Lemma 2.1 that the invariant set Λ is uniformly hyperbolic. This completes the proof. □

Therefore, by Lemmas 2.2, 3.4, and 3.5, one has that Theorem 3.2 holds.

Theorem 3.3. For system (3.4), suppose that $p(x) = \prod_{i=1}^{m}(x-\alpha_i)$, where $\alpha_1 < \alpha_2 < \cdots < \alpha_m$ are real numbers, and $m \geq 2$. For any fixed a_3, b_1, and c_2 with $|c_2| < 1$, and sufficiently large $|a_1|$, there exist a Smale horseshoe and a hyperbolic invariant set on which F is topologically conjugate to the two-sided fullshift on m symbols. Consequently, F is chaotic in the sense of both Li-Yorke and Devaney.

Proof. Denote $\lambda := \frac{1}{|c_2|}$. By the properties of $p(x)$, there are positive constants η_i, $1 \leq i \leq m$, such that the sets $V_i = [\alpha_i - \eta_i, \alpha_i + \eta_i]$, $1 \leq i \leq m$, are pairwise disjoint, and $|p'(x)| > 0$ for all $x \in V_i$, $1 \leq i \leq m$.

Set

$$U := [\alpha_1 - \eta_1, \alpha_m + \eta_m] \times [b_1(\alpha_1 - \eta_1), b_1(\alpha_m + \eta_m)] \times [c_2 b_1(\alpha_1 - \eta_1), c_2 b_1(\alpha_m + \eta_m)],$$

and

$$\Lambda := \bigcap_{-\infty}^{+\infty} F^j(U).$$

Set

$$M_0 := \inf_{x \in \cup_{i=1}^{m} V_i} |p'(x)|, \quad M_1 := \min_{1 \leq i \leq m} |p(\alpha_i \pm \eta_i)|;$$

$$N_0 := \max\left\{ \frac{\lambda + |a_2| + |a_3|}{M_0}, \ \frac{\lambda |a_3 b_1 c_2| + |b_1 c_2| + |a_2 b_1|}{M_0 |c_2|} \right\}.$$

$$N_1 := \frac{(1 + |a_2 b_1| + |a_3 c_2 b_1|) \max\{|\alpha_1 - \eta_1|, |\alpha_n + \eta_n|\}}{M_1}.$$

If $|a_1| > N_0$, by applying similar discussions in the proof of Lemma 3.3, one has that the invariant set Λ is uniformly hyperbolic, where the constant N_0 is similar with the constant specified in (3.9).

If $|a_1| > N_1$, by applying simple calculation and the fact that $p(x)$ has simple real roots, one has that $F(U) \cap U$ has m connected components.

Hence, if $|a_1| > \max\{N_0, N_1\}$, then there exist a Smale horseshoe and a hyperbolic invariant set on which F is topologically conjugate to the two-sided fullshift on m symbols. This, together with Lemma 2.2, yields that F is chaotic in the sense of both Li-Yorke and Devaney. This completes the proof. ☐

3.2 The maps with the dimension of the unstable subspace equals to two

In this subsection, we consider a subclass of the maps (1.3) satisfying that $a_3 = c_2 = c_3 = 0$, $a_2b_3 \neq 0$, $c_1 = 1$, and $a_2b_3 \neq 0$, which are represented as follows: $F : (x,y,z) \in \mathbb{R}^3 \to (f_1, f_2, f_3) \in \mathbb{R}^3$:

$$\left\{\begin{array}{l} f_1(x,y,z) = a_1p(x) + a_2y \\ f_2(x,y,z) = b_1x + b_2q(y) + b_3z \\ f_3(x,y,z) = x \end{array}\right. . \tag{3.17}$$

The derivative of F and the determinant are

$$DF = \begin{pmatrix} a_1p'(x) & a_2 & 0 \\ b_1 & b_2q'(y) & b_3 \\ 1 & 0 & 0 \end{pmatrix} \tag{3.18}$$

and $a_2b_3 \neq 0$, respectively. The inverse of F is $H : (x,y,z) \in \mathbb{R}^3 \to (h_1, h_2, h_3) \in \mathbb{R}^3$:

$$\left\{\begin{array}{l} h_1(x,y,z) = z \\ h_2(x,y,z) = \frac{1}{a_2}(x - a_1p(z)) \\ h_3(x,y,z) = \frac{1}{b_3}(y - b_1z - b_2q(\frac{1}{a_2}(x - a_1p(z)))) \end{array}\right. \tag{3.19}$$

and the derivative of H is

$$DF^{-1} = \begin{pmatrix} 0 & 0 & 1 \\ \frac{1}{a_2} & 0 & -\frac{a_1}{a_2}p'(z) \\ -\frac{b_2}{a_2b_3}q'(\frac{1}{a_2}(x - a_1p(z))) & \frac{1}{b_3} & -\frac{b_1}{b_3} + \frac{a_1b_2}{a_2b_3}q'(\frac{1}{a_2}(x - a_1p(z)))p'(z) \end{pmatrix}. \tag{3.20}$$

The determinant is $1/a_2b_3$. Hence, polynomial maps (3.17) are diffeomorphisms with constant Jacobian and polynomial inverse.

To begin our study, a toy model is introduced as follows:

$$\left\{\begin{array}{l} f_1(x,y,z) = a_1x(1-x) + a_2y \\ f_2(x,y,z) = b_1x + b_2y(1-y) + b_3z \\ f_3(x,y,z) = x \end{array}\right. .$$

This model can be thought of as another generalization of the Hénon map in three-dimensional spaces.

Now, we show that for certain parameters, there exist uniformly hyperbolic invariant sets and chaotic dynamics.

Theorem 3.4. Suppose that there are $1 \leq i_0 < r_1$ and $1 \leq j_0 < r_2$ such that $0 = \alpha_{i_0} < \alpha_{i_0+1}$, $0 = \xi_{j_0} < \xi_{j_0+1}$, and $m_{i_0} = m_{i_0+1} = n_{j_0} = n_{j_0+1} = 1$, that is, there are two distinct non-negative zeros of $p(x)$ and $q(y)$. If

$$a_2 < 0 \text{ and } a_1p'(0) > 0, \text{ and } b_1 \leq 0, \ b_3 < 0, \text{ and } b_2q'(0) > 0, \tag{3.21}$$

then, for fixed a_2, b_1, and b_3, and sufficiently large $|a_1|$ and $|b_2|$, there are a Smale horseshoe for the map (3.17), and a uniformly hyperbolic invariant set Λ on which F is topologically conjugate to two-sided fullshift on four symbols. Therefore, they are Li-Yorke chaotic as well as Devaney chaotic.

Consider the compact set

$$U := [0, \alpha_{i_0+1}] \times [0, \xi_{j_0+1}] \times [0, \alpha_{i_0+1}].$$

Take a constant $\lambda > 1$. The invariant set is denoted by

$$\Lambda := \bigcap_{-\infty}^{+\infty} F^i(U).$$

Lemma 3.6. Under the assumptions of Theorem 3.4, one has that $F(U) \cap U$ and $F^{-1}(U) \cap U$ have four non-empty connected components, respectively.

Proof. It is to show that $F(U) \cap U$ has four connected components. We will describe how to think of the image of $F(U)$.

First, it is to find the position of the image of the vertices of the cubic U, that is, the image of $A, B, C, D, A', B', C', D'$ under F. We only need to know the relative position of $F(A)$, $F(B)$, $F(C)$, $F(D)$, $F(A')$, $F(B')$, $F(C')$, $F(D')$. The vertices of the cubic U are:

$$A = (\alpha_{i_0+1}, 0, 0), B = (\alpha_{i_0+1}, \xi_{j_0+1}, 0), C = (0, \xi_{j_0+1}, 0), D = (0, 0, 0),$$

$$A' = (\alpha_{i_0+1}, 0, \alpha_{i_0+1}), B' = (\alpha_{i_0+1}, \xi_{j_0+1}, \alpha_{i_0+1}), C' = (0, \xi_{j_0+1}, \alpha_{i_0+1}), D' = (0, 0, \alpha_{i_0+1});$$

the image of the vertices under F are

$$F(A) = (0, b_1\alpha_{i_0+1}, \alpha_{i_0+1}),\ F(B) = (a_2\xi_{j_0+1}, b_1\alpha_{i_0+1}, \alpha_{i_0+1}), F(C) = (a_2\xi_{j_0+1}, 0, 0),$$

$$F(D) = (0, 0, 0), F(A') = (0, (b_1 + b_3)\alpha_{i_0+1}, \alpha_{i_0+1}),\ F(B') = (a_2\xi_{j_0+1}, (b_1 + b_3)\alpha_{i_0+1}, \alpha_{i_0+1}),$$

$$F(C') = (a_2\xi_{j_0+1}, b_3\alpha_{i_0+1}, 0),\ F(D') = (0, b_3\alpha_{i_0+1}, 0).$$

Set

$$AD := \{(x, y, z) :\ x \in [0, \alpha_{i_0+1}], y = 0, z = 0\},$$

$$AB := \{(x, y, z) :\ x = \alpha_{i_0+1}, y \in [0, \xi_{j_0+1}], z = 0\},$$

$$AA' := \{(x,y,z):\ x = \alpha_{i_0+1}, y = 0, z \in [0, \alpha_{i_0+1}]\},$$

$$ABCD := \{(x,y,z):\ (x,y) \in [0,\alpha_{i_0+1}] \times [0, \xi_{j_0+1}], z = 0\}.$$

Second, it is to determine the image of the plane $ABCD$ by using the expression (3.17). By (3.17), $F(AD)$ is a parabola along the x-axis, and $F(AB)$ is a parabola along y-axis. The image of $ABCD$ under F can be thought of as a movement of the parabola $F(AD)$ along another parabola $F(AB)$, but the direction of $F(AD)$ should not change too much.

Finally, it is to study the graph of $F(U)$. By (3.17), $F(AA')$ is a line segment. We push forward the surface $F(ABCD)$ obtained in the previous step along $F(AA')$. The graph of $F(U)$ comes out!

Hence, there are four non-empty connected components of $F(U) \cap U$ for sufficiently large $|a_1|$ and $|b_2|$.

Now, it is to study $F^{-1}(U) \cap U$. The image of the vertices of U under $F^{-1} = H$ are

$$H(A) = \left(0, \frac{\alpha_{i_0+1}}{a_2}, \frac{-b_2 q(\alpha_{i_0+1}/a_2)}{b_3}\right), H(B) = \left(0, \frac{\alpha_{i_0+1}}{a_2}, \frac{(\xi_{j_0+1} - b_2 q(\alpha_{i_0+1}/a_2))}{b_3}\right),$$

$$H(C) = (0, 0, \xi_{j_0+1}/b_3), H(D) = (0,0,0),$$

$$H(A') = \left(\alpha_{i_0+1}, \frac{\alpha_{i_0+1}}{a_2}, \frac{(-b_1\alpha_{i_0+1} - b_2 q(\alpha_{i_0+1}/a_2))}{b_3}\right),$$

$$H(B') = \left(\alpha_{i_0+1}, \frac{\alpha_{i_0+1}}{a_2}, \frac{(\xi_{j_0+1} - b_1\alpha_{i_0+1} - b_2 q(\alpha_{i_0+1}/a_2))}{b_3}\right),$$

$$H(C') = \left(\alpha_{i_0+1}, 0, \frac{(\xi_{j_0+1} - b_1\alpha_{i_0+1})}{b_3}\right), H(D') = \left(\alpha_{i_0+1}, 0, \frac{-b_1\alpha_{i_0+1}}{b_3}\right).$$

By (3.19), one has that for fixed a_2 and b_3, and sufficiently large $|a_1|$ and $|b_2|$, given any $(x_0, y_0, z_0) \in U$,

$$\begin{cases} x_0 = z \\ y_0 = \frac{1}{a_2}(x - a_1 p(z)) \\ z_0 = \frac{1}{b_3}(y - b_1 z - b_2 q(\frac{1}{a_2}(x - a_1 p(z)))) \end{cases},$$

if there is a solution of the above equations, then there should exist four different solutions by Lemma 3.1. Further, it follows from $F^{-1}(U) \cap U = F^{-1}(U \cap F(U))$, the fact that F is a diffeomorphism and the above geometric description of $F(U)$ that $F^{-1}(U) \cap U$ has four non-empty connected components. The proof is completed. □

Lemma 3.7. Under the assumptions of Theorem 3.4, the invariant set Λ is uniformly hyperbolic.

Proof. Fix a_2, b_1, and b_3, it follows from (i) and (ii) of Lemma 3.1 that we could assume $|a_1|$ and $|b_2|$ are sufficiently large such that for $w = (x, y, z) \in U \cap F^{-1}(U)$, one has that $x \in [\alpha_{i_0}, \delta_1] \cup [\delta_1', \alpha_{i_0+1}]$ and $y \in [\xi_{j_0}, \delta_2] \cup [\delta_2', \xi_{j_0+1}]$. For any point $w_0 = (x_0, y_0, z_0) \in U$, set $w_1 = F(w_0) = (x_1, y_1, z_1)$ and $w_{-1} = F^{-1}(w_0) = (x_{-1}, y_{-1}, z_{-1})$.

Set

$$M_0 := \min\left\{ \min_{x\in[0,\delta_1]\cup[\delta_1',\alpha_{i_0+1}]} |p'(x)|, \min_{y\in[0,\delta_2]\cup[\delta_2',\xi_{j_0+1}]} |q'(y)| \right\}.$$

Suppose that

$$|a_1| \geq \frac{\max\{\lambda + |a_2|, 1 + |a_2 b_3|\lambda + |a_2|(1 + |b_1|)\}}{M_0}, \ |b_2| \geq \frac{\lambda + |b_1| + |b_3|}{M_0}. \tag{3.22}$$

Now, it is to show that for any point $w_0 \in U \cap F^{-1}(U)$, $m_{\mathcal{C},w_0} \geq \lambda$.

Introduce the unit cone

$$K_1(\mathbb{R}^2, \mathbb{R}) = \left\{ \mathbf{v} = \begin{pmatrix} v_0 \\ u_0 \end{pmatrix} : \ v_0 = (v_{01}, v_{02})^T \in \mathbb{R}^2, \ u_0 \in \mathbb{R}, \text{ and } |u_0| \leq |v_0| \right\},$$

where $|v_0| = \max\{|v_{01}|, |v_{02}|\}$ and $|\mathbf{v}| = \max\{|v_0|, |u_0|\}$.

For any $\begin{pmatrix} v_0 \\ u_0 \end{pmatrix} \in K_1(\mathbb{R}^2, \mathbb{R})$, $\begin{pmatrix} v_1 \\ u_1 \end{pmatrix} = DF_{w_0} \begin{pmatrix} v_0 \\ u_0 \end{pmatrix}$. By (3.18),

$$\begin{cases} v_{11} = a_1 p'(x_0) v_{01} + a_2 v_{02} \\ v_{12} = b_1 v_{01} + b_2 q'(y_0) v_{02} + b_3 u_0 \\ u_1 = v_{01} \end{cases}.$$

By (3.22), if $|v_0| = |v_{01}|$, then

$$|v_{11}| \geq |v_{01}||a_1 p'(x_0)| - |a_2||v_{01}| \geq \lambda |v_{01}|;$$

if $|v_0| = |v_{02}|$, then

$$|v_{12}| \geq |v_{02}||b_2 q'(y_0)| - |b_1||v_{02}| - |b_3||u_0| \geq \lambda |v_{02}|,$$

which implies that $|(v_1, u_1)| \geq \lambda |(v_0, u_0)|$. Hence, $m_{\mathcal{C},w_0} \geq \lambda$.

Next, it is to show that $m'_{\mathcal{C},w_0} \geq \lambda$. for any point $w_0 \in U \cap F(U)$, take any $\begin{pmatrix} v_0 \\ u_0 \end{pmatrix} \notin K_1(\mathbb{R}^2, \mathbb{R})$, that is , $|u_0| > |v_0|$, suppose $\begin{pmatrix} v_{-1} \\ u_{-1} \end{pmatrix} = DF^{-1}_{w_0} \begin{pmatrix} v_0 \\ u_0 \end{pmatrix}$, by (3.20), one has

$$\begin{cases} v_{-11} = u_0 \\ v_{-12} = \frac{1}{a_2} v_{01} - \frac{a_1}{a_2} p'(z_0) u_0 \\ u_{-1} = -\frac{b_2}{a_2 b_3} q'(\frac{1}{a_2}(x_0 - a_1 p(z_0))) v_{01} + \frac{1}{b_3} v_{02} - \frac{b_1}{b_3} u_0 + \frac{a_1 b_2}{a_2 b_3} q'(\frac{1}{a_2}(x_0 - a_1 p(z_0))) p'(z_0) u_0 \end{cases}.$$

So, by (3.22),

$$\begin{aligned}|u_{-1}| \geq & \left|\frac{a_1 b_2}{a_2 b_3} q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_0))\right) p'(z_0)\right| |u_0| - \left|\frac{b_2}{a_2 b_3} q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_0))\right)\right| |v_{01}| - \frac{|v_{02}|}{|b_3|} - \frac{|b_1|}{|b_3|}|u_0| \\ \geq & \left|\frac{a_1 b_2}{a_2 b_3} q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_1))\right) p'(z_0)\right| |u_0| - \left|\frac{b_2}{a_2 b_3} q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_0))\right)\right| |v_0| - \frac{|v_{02}|}{|b_3|} - \frac{|b_1|}{|b_3|}|u_0| \\ \geq & \left|\frac{a_1 b_2}{a_2 b_3} q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_0))\right) p'(z_0)\right| |u_0| - \left|\frac{b_2}{a_2 b_3} q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_0))\right)\right| |u_0| - \frac{|u_0|}{|b_3|} - \frac{|b_1|}{|b_3|}|u_0| \\ \geq & \lambda |u_0|,\end{aligned}$$

which yields that $m'_{\mathcal{C},w_0} \geq \lambda$.

It follows from Lemma 2.1 that the invariant set Λ is uniformly hyperbolic. This completes the proof. ☐

Therefore, it follows from Lemmas 2.2, 3.6, and 3.7 that Theorem 3.4 holds.

Theorem 3.5. Suppose that there are $1 \leq i_0 < r_1$ and $1 \leq j_0 < r_2$ such that $0 < \alpha_{i_0} < \alpha_{i_0+1}$ and $0 < \xi_{j_0} < \xi_{j_0+1}$, that is, there are two distinct positive zeros of $p(x)$ and $q(y)$. Set

$$M_1 := \max\{a_2\xi_{j_0}, a_2\xi_{j_0+1}\},\ M_2 := \min\{a_2\xi_{j_0}, a_2\xi_{j_0+1}\},$$

$$N_1 := \max\{b_1\alpha_{i_0} + b_3\alpha_{i_0}, b_1\alpha_{i_0} + b_3\alpha_{i_0+1}, b_1\alpha_{i_0+1} + b_3\alpha_{i_0}, b_1\alpha_{i_0+1} + b_3\alpha_{i_0+1}\},$$

$$N_2 := \min\{b_1\alpha_{i_0} + b_3\alpha_{i_0}, b_1\alpha_{i_0} + b_3\alpha_{i_0+1}, b_1\alpha_{i_0+1} + b_3\alpha_{i_0}, b_1\alpha_{i_0+1} + b_3\alpha_{i_0+1}\}.$$

Consider the following different situations:

(i). $M_1 < \alpha_{i_0}$ and $a_1 p^{(m_{i_0})}(\alpha_{i_0}) > 0$, and $N_1 < \xi_{j_0}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) > 0$;

(ii). $M_1 < \alpha_{i_0}$ and $a_1 p^{(m_{i_0})}(\alpha_{i_0}) > 0$, and $N_2 > \xi_{j_0+1}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) < 0$;

(iii). $M_2 > \alpha_{i_0+1}$ and $a_1 p^{(m_{i_0})}(\alpha_{i_0}) < 0$, and $N_1 < \xi_{j_0}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) > 0$;

(iv). $M_2 > \alpha_{i_0+1}$ and $a_1 p^{(m_{i_0})}(\alpha_{i_0}) < 0$, and $N_2 > \xi_{j_0+1}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) < 0$.

Then, for fixed a_2, b_1, and b_3, and sufficiently large $|a_1|$ and $|b_2|$, there exists a Smale horseshoe for the map (3.17), especially, there is a uniformly hyperbolic invariant set Λ on which F is topologically conjugate to two-sided fullshift on four symbols. Therefore, they are Li-Yorke chaotic as well as Devaney chaotic.

Remark 3.1. The four different situations in the assumptions of Theorem 3.5 is the generalization of the assumptions (3.10) in Theorem 3.2.

Consider the compact set

$$U := [\alpha_{i_0}, \alpha_{i_0+1}] \times [\xi_{j_0}, \xi_{j_0+1}] \times [\alpha_{i_0}, \alpha_{i_0+1}].$$

Fix a constant $\lambda > 1$. The invariant set is denoted by

$$\Lambda := \bigcap_{-\infty}^{+\infty} F^i(U).$$

Lemma 3.8. In Cases (i)-(iv) of Theorem 3.5, for fixed a_2, b_1, and b_3, and sufficiently large $|a_1|$ and $|b_2|$, one has that $F(U) \cap U$ and $F^{-1}(U) \cap U$ have four non-empty connected components, respectively.

Proof. It is to show that $F(U) \cap U$ has four connected components under the assumptions of Theorem 3.5. We will give the geometric description of $F(U)$ in the case: $a_2 > 0$, $b_3 > 0$, $b_1 = 0$, $a_2\xi_{j_0+1} < \alpha_{i_0}$, $b_3\alpha_{i_0+1} < \xi_{j_0}$, $a_1 p^{(m_{i_0})}(\alpha_{i_0}) > 0$, and $b_2 q^{(n_{j_0})}(\xi_{j_0}) > 0$. The graph for other parameters could be obtained similarly.

First, Since U is a cubic, the first step is to find the position of the vertices, that is, the image of $A, B, C, D, A', B', C', D'$. We should know the relative position of $F(A)$, $F(B)$, $F(C)$, $F(D)$, $F(A')$, $F(B')$, $F(C')$, $F(D')$. The vertices of the cubic U are listed as follows:

$$A = (\alpha_{i_0+1}, \xi_{j_0}, \alpha_{i_0}), B = (\alpha_{i_0+1}, \xi_{j_0+1}, \alpha_{i_0}), C = (\alpha_{i_0}, \xi_{j_0+1}, \alpha_{i_0}), D = (\alpha_{i_0}, \xi_{j_0}, \alpha_{i_0}),$$

$$A' = (\alpha_{i_0+1}, \xi_{j_0}, \alpha_{i_0+1}), B' = (\alpha_{i_0+1}, \xi_{j_0+1}, \alpha_{i_0+1}), C' = (\alpha_{i_0}, \xi_{j_0+1}, \alpha_{i_0+1}), D' = (\alpha_{i_0}, \xi_{j_0}, \alpha_{i_0+1});$$

the image of these points under F are

$$F(A) = (a_2\xi_{j_0}, b_1\alpha_{i_0+1} + b_3\alpha_{i_0}, \alpha_{i_0+1}),\ F(B) = (a_2\xi_{j_0+1}, b_1\alpha_{i_0+1} + b_3\alpha_{i_0}, \alpha_{i_0+1}),$$

$$F(C) = (a_2\xi_{j_0+1}, b_1\alpha_{i_0} + b_3\alpha_{i_0}, \alpha_{i_0}),\ F(D) = (a_2\xi_{j_0}, b_1\alpha_{i_0} + b_3\alpha_{i_0}, \alpha_{i_0}),$$

$$F(A') = (a_2\xi_{j_0}, b_1\alpha_{i_0+1} + b_3\alpha_{i_0+1}, \alpha_{i_0+1}),\ F(B') = (a_2\xi_{j_0+1}, b_1\alpha_{i_0+1} + b_3\alpha_{i_0+1}, \alpha_{i_0+1}),$$

$$F(C') = (a_2\xi_{j_0+1}, b_1\alpha_{i_0} + b_3\alpha_{i_0+1}, \alpha_{i_0}),\ F(D') = (a_2\xi_{j_0}, b_1\alpha_{i_0} + b_3\alpha_{i_0+1}, \alpha_{i_0}).$$

It is evident that the image of the vertices are not contained in U.

Set

$$AD := \{(x,y,z):\ x\in[\alpha_{i_0},\alpha_{i_0+1}], y=\xi_{j_0}, z=\alpha_{i_0}\},$$

$$AB := \{(x,y,z):\ x=\alpha_{i_0+1}, y\in[\xi_{j_0},\xi_{j_0+1}], z=\alpha_{i_0}\},$$

$$AA' := \{(x,y,z):\ x=\alpha_{i_0+1}, y=\xi_{j_0}, z\in[\alpha_{i_0},\alpha_{i_0+1}]\},$$

$$ABCD := \{(x,y,z):\ (x,y)\in[\alpha_{i_0},\alpha_{i_0+1}]\times[\xi_{j_0},\xi_{j_0+1}], z=\alpha_{i_0}\}.$$

Second, it is to determine the image of the plane $ABCD$ under the map F. By (3.17), $F(AD)$ is a parabola along the x-axis, and $F(AB)$ is a parabola along y-axis. The image $F(ABCD)$ can be regarded as a movement of the parabola $F(AD)$ along another parabola $F(AB)$, but the direction of $F(AD)$ should not vary too much.

Finally, it is to investigate the graph of $F(U)$. It follows from (3.17) that $F(AA')$ is a line segment. The graph of $F(U)$ can be obtained by moving the surface $F(ABCD)$ obtained in the previous step along $F(AA')$.

Hence, there are four non-empty connected components of $F(U)\cap U$ for sufficiently large $|a_1|$ and $|b_2|$.

Now, it is to study $F^{-1}(U)\cap U$. The image of the vertices of U under $F^{-1}=H$ are

$$H(A)=\left(\alpha_{i_0}, \frac{\alpha_{i_0+1}}{a_2}, \frac{(\xi_{j_0}-b_1\alpha_{i_0}-b_2q(\alpha_{i_0+1}/a_2))}{b_3}\right),$$

$$H(B)=\left(\alpha_{i_0}, \frac{\alpha_{i_0+1}}{a_2}, \frac{(\xi_{j_0+1}-b_1\alpha_{i_0}-b_2q(\alpha_{i_0+1}/a_2))}{b_3}\right),$$

$$H(C)=\left(\alpha_{i_0}, \frac{\alpha_{i_0}}{a_2}, \frac{(\xi_{j_0+1}-b_1\alpha_{i_0}-b_2q(\alpha_{i_0}/a_2))}{b_3}\right),$$

$$H(D)=\left(\alpha_{i_0}, \frac{\alpha_{i_0}}{a_2}, \frac{(\xi_{j_0}-b_1\alpha_{i_0}-b_2q(\alpha_{i_0}/a_2))}{b_3}\right),$$

$$H(A')=\left(\alpha_{i_0+1}, \frac{\alpha_{i_0+1}}{a_2}, \frac{(\xi_{j_0}-b_1\alpha_{i_0+1}-b_2q(\alpha_{i_0+1}/a_2))}{b_3}\right),$$

$$H(B')=\left(\alpha_{i_0+1}, \frac{\alpha_{i_0+1}}{a_2}, \frac{(\xi_{j_0+1}-b_1\alpha_{i_0+1}-b_2q(\alpha_{i_0+1}/a_2))}{b_3}\right),$$

$$H(C')=\left(\alpha_{i_0+1}, \frac{\alpha_{i_0}}{a_2}, \frac{(\xi_{j_0+1}-b_1\alpha_{i_0+1}-b_2q(\alpha_{i_0}/a_2))}{b_3}\right),$$

$$H(D')=\left(\alpha_{i_0+1}, \frac{\alpha_{i_0}}{a_2}, \frac{(\xi_{j_0}-b_1\alpha_{i_0+1}-b_2q(\alpha_{i_0}/a_2))}{b_3}\right).$$

By (3.19), given any $(x_0, y_0, z_0) \in U$,

$$\begin{cases} x_0 = z \\ y_0 = \frac{1}{a_2}(x - a_1 p(z)) \\ z_0 = \frac{1}{b_3}(y - b_1 z - b_2 q(\frac{1}{a_2}(x - a_1 p(z)))) \end{cases},$$

there should exist four different solutions by Lemma 3.1 if the solution set is non-empty. Further, since F is a diffeomorphism, $F^{-1}(U) \cap U = F^{-1}(U \cap F(U))$. This, together with the above geometric description of $F(U)$, yields that $F^{-1}(U) \cap U$ has four non-empty connected components.

This completes the proof. □

Remark 3.2. In the last section, we provide an example with the help of Mathematica software to draw the graph of $F(U) \cap U$ and $F^{-1}(U) \cap U$ to illustrate the results of Theorem 3.5.

Lemma 3.9. In Cases (i)-(iv) of Theorem 3.5, for fixed a_2, b_1, and b_3, and sufficiently large $|a_1|$ and $|b_2|$, one has that the invariant set is uniformly hyperbolic.

Proof. Without loss of generality, assume that a_2, b_1, and b_3 are fixed, $|a_1|$ and $|b_2|$ are large enough. By (i) and (ii) of Lemma 3.1, we could assume that $|a_1|$ and $|b_2|$ are sufficiently large such that for $w = (x, y, z) \in U \cap F^{-1}(U)$, one has that $x \in (\alpha_{i_0}, \delta_1] \cup [\delta_1', \alpha_{i_0+1})$ and $y \in (\xi_{j_0}, \delta_2] \cup [\delta_2', \xi_{j_0+1})$. For any point $w_0 = (x_0, y_0, z_0) \in U$, denote $w_1 = F(w_0) = (x_1, y_1, z_1)$ and $w_{-1} = F^{-1}(w_0) = (x_{-1}, y_{-1}, z_{-1})$.

From (3.1), (3.2), and Lemma 3.1, it follows that for any $w = (x, y, z) \in U \cap F^{-1}(U)$,

$$\text{Case (i)}: \frac{|a_1 p'(x)|}{|a_1 p(x)|} \geq \frac{\max\{\lambda + |a_2|, 1 + |a_2 b_3|\lambda + |a_2|(1 + |b_1|)\}}{\alpha_{i_0} - M_1}, \frac{|b_2 q'(y)|}{|b_2 q(y)|} \geq \frac{\lambda + |b_1| + |b_3|}{\xi_{j_0} - N_1}; \tag{3.23}$$

$$\text{Case (ii)}: \frac{|a_1 p'(x)|}{|a_1 p(x)|} \geq \frac{\max\{\lambda + |a_2|, 1 + |a_2 b_3|\lambda + |a_2|(1 + |b_1|)\}}{\alpha_{i_0} - M_1}, \frac{|b_2 q'(y)|}{|b_2 q(y)|} \geq \frac{\lambda + |b_1| + |b_3|}{N_2 - \xi_{j_0+1}}; \tag{3.24}$$

$$\text{Case (iii)}: \frac{|a_1 p'(x)|}{|a_1 p(x)|} \geq \frac{\max\{\lambda + |a_2|, 1 + |a_2 b_3|\lambda + |a_2|(1 + |b_1|)\}}{M_2 - \alpha_{i_0+1}}, \frac{|b_2 q'(y)|}{|b_2 q(y)|} \geq \frac{\lambda + |b_1| + |b_3|}{\xi_{j_0} - N_1}; \tag{3.25}$$

$$\text{Case (iv)}: \frac{|a_1 p'(x)|}{|a_1 p(x)|} \geq \frac{\max\{\lambda + |a_2|, 1 + |a_2 b_3|\lambda + |a_2|(1 + |b_1|)\}}{M_2 - \alpha_{i_0+1}}, \frac{|b_2 q'(y)|}{|b_2 q(y)|} \geq \frac{\lambda + |b_1| + |b_3|}{N_2 - \xi_{j_0+1}}. \tag{3.26}$$

Consider the point $w_0 \in U \cap F^{-1}(U)$, by (3.17),

$$\left\{ \begin{array}{l} x_1 = a_1 p(x_0) + a_2 y_0 \\ y_1 = b_1 x_0 + b_2 q(y_0) + b_3 z_0 \\ z_1 = x_0 \end{array} \right. .$$

So, $a_1 p(x_0) = x_1 - a_2 y_0$ and $b_2 q(y_0) = y_1 - b_1 x_0 - b_3 z_0$. Hence, one has

$$\text{Case (i)}: \quad a_1 p(x_0) \geq \alpha_{i_0} - M_1,\ b_2 q(y_0) \geq \xi_{j_0} - N_1; \tag{3.27}$$

$$\text{Case (ii)}: \quad a_1 p(x_0) \geq \alpha_{i_0} - M_1,\ -b_2 q(y_0) \geq N_2 - \xi_{j_0+1}; \tag{3.28}$$

$$\text{Case (iii)}: \quad -a_1 p(x_0) \geq M_2 - \alpha_{i_0+1},\ b_2 q(y_0) \geq \xi_{j_0} - N_1; \tag{3.29}$$

$$\text{Case (iv)}: \quad -a_1 p(x_0) \geq M_2 - \alpha_{i_0+1},\ -b_2 q(y_0) \geq N_2 - \xi_{j_0+1}. \tag{3.30}$$

For any point $w_0 \in U \cap F(U)$, it follows from (3.19) that

$$\left\{ \begin{array}{l} x_{-1} = z_0 \\ y_{-1} = \frac{1}{a_2}(x_0 - a_1 p(z_0)) \\ z_{-1} = \frac{1}{b_3}(y_0 - b_1 z_0 - b_2 q(\frac{1}{a_2}(x_0 - a_1 p(z_0)))) \end{array} \right. .$$

So, $a_1 p(x_{-1}) = a_1 p(z_0) = x_0 - a_2 y_{-1}$ and $b_2 q(y_{-1}) = b_2 q(\frac{1}{a_2}(x_0 - a_1 p(z_0))) = y_0 - b_1 z_0 - b_3 z_{-1}$. Thus, one has

$$\text{Case (i)}: \quad a_1 p(x_{-1}) \geq \alpha_{i_0} - M_1,\ b_2 q(y_{-1}) \geq \xi_{j_0} - N_1; \tag{3.31}$$

$$\text{Case (ii)}: \quad a_1 p(x_{-1}) \geq \alpha_{i_0} - M_1,\ -b_2 q(y_{-1}) \geq N_2 - \xi_{j_0+1}; \tag{3.32}$$

$$\text{Case (iii)}: \quad -a_1 p(x_{-1}) \geq M_2 - \alpha_{i_0+1},\ b_2 q(y_{-1}) \geq \xi_{j_0} - N_1; \tag{3.33}$$

$$\text{Case (iv)}: \quad -a_1 p(x_{-1}) \geq M_2 - \alpha_{i_0+1},\ -b_2 q(y_{-1}) \geq N_2 - \xi_{j_0+1}. \tag{3.34}$$

By (3.23)–(3.26) and (3.27)–(3.30), one has that for any point $w_0 \in U \cap F^{-1}(U)$,

$$\text{Cases (i)–(iv)}: \ |a_1 p'(x_0)| \geq \lambda + |a_2|,\ |b_2 q'(y_0)| \geq \lambda + |b_1| + |b_3|. \tag{3.35}$$

Now, it is to show that for any point $w_0 \in U \cap F^{-1}(U)$, $m_{\mathcal{C},w_0} \geq \lambda > 1$.

Consider the unit cone

$$K_1(\mathbb{R}^2, \mathbb{R}) = \left\{ \mathbf{v} = \begin{pmatrix} v_0 \\ u_0 \end{pmatrix} : \ v_0 = (v_{01}, v_{02})^T \in \mathbb{R}^2,\ u_0 \in \mathbb{R},\ \text{and } |u_0| \leq |v_0| \right\},$$

where $|v_0| = \max\{|v_{01}|, |v_{02}|\}$ and $|\mathbf{v}| = \max\{|v_0|, |u_0|\}$.

Suppose that $\begin{pmatrix} v_1 \\ u_1 \end{pmatrix} = DF_{w_0} \begin{pmatrix} v_0 \\ u_0 \end{pmatrix}$, where $\begin{pmatrix} v_0 \\ u_0 \end{pmatrix} \in K_1(\mathbb{R}^2, \mathbb{R})$. By (3.18),

$$\left\{ \begin{array}{l} v_{11} = a_1 p'(x_0) v_{01} + a_2 v_{02} \\ v_{12} = b_1 v_{01} + b_2 q'(y_0) v_{02} + b_3 u_0 \\ u_1 = v_{01} \end{array} \right. .$$

By (3.35), if $|v_0| = |v_{01}|$, then

$$|v_{11}| \geq |v_{01}||a_1 p'(x_0)| - |a_2||v_{01}| \geq \lambda |v_{01}|;$$

if $|v_0| = |v_{02}|$, then

$$|v_{12}| \geq |v_{02}||b_2 q'(y_0)| - |b_1||v_{02}| - |b_3||u_0| \geq \lambda |v_{02}|,$$

which implies that $|(v_1, u_1)| \geq \lambda |(v_0, u_0)|$. Hence, $m_{\mathcal{C},w_0} \geq \lambda > 1$.

By (3.23)–(3.26) and (3.31)–(3.34), one has that for any point $w_0 \in U \cap F(U)$,

$$|a_1 p'(x_{-1})| = |a_1 p'(z_0)| \geq 1 + |a_2 b_3|\lambda + |a_2|(1 + |b_1|),\ |b_2 q'(y_{-1})| = \left| b_2 q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_0))\right) \right| \geq 1. \tag{3.36}$$

Next, it is to show that then $m'_{\mathcal{C},w_0} \geq \lambda > 1$. For any $\begin{pmatrix} v_0 \\ u_0 \end{pmatrix} \notin K_1(\mathbb{R}^2, \mathbb{R})$, that is , $|u_0| > |v_0|$, suppose $\begin{pmatrix} v_{-1} \\ u_{-1} \end{pmatrix} = DF_{w_0}^{-1} \begin{pmatrix} v_0 \\ u_0 \end{pmatrix}$, by (3.20), one has

$$\left\{ \begin{array}{l} v_{-11} = u_0 \\ v_{-12} = \frac{1}{a_2} v_{01} - \frac{a_1}{a_2} p'(z_0) u_0 \\ u_{-1} = -\frac{b_2}{a_2 b_3} q'(\frac{1}{a_2}(x_0 - a_1 p(z_0))) v_{01} + \frac{1}{b_3} v_{02} - \frac{b_1}{b_3} u_0 + \frac{a_1 b_2}{a_2 b_3} q'(\frac{1}{a_2}(x_0 - a_1 p(z_0))) p'(z_0) u_0 \end{array} \right. .$$

So, by (3.36),

$$\begin{aligned} |u_{-1}| \geq & \left| \frac{a_1 b_2}{a_2 b_3} q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_0))\right) p'(z_0) \right| |u_0| - \left| \frac{b_2}{a_2 b_3} q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_0))\right) \right| |v_{01}| - \frac{|v_{02}|}{|b_3|} - \frac{|b_1|}{|b_3|} |u_0| \\ \geq & \left| \frac{a_1 b_2}{a_2 b_3} q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_1))\right) p'(z_0) \right| |u_0| - \left| \frac{b_2}{a_2 b_3} q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_0))\right) \right| |v_0| - \frac{|v_{02}|}{|b_3|} - \frac{|b_1|}{|b_3|} |u_0| \\ \geq & \left| \frac{a_1 b_2}{a_2 b_3} q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_0))\right) p'(z_0) \right| |u_0| - \left| \frac{b_2}{a_2 b_3} q'\left(\frac{1}{a_2}(x_0 - a_1 p(z_0))\right) \right| |u_0| - \frac{|u_0|}{|b_3|} - \frac{|b_1|}{|b_3|} |u_0| \\ \geq & \lambda |u_0|, \end{aligned}$$

which yields that $m'_{\mathcal{C},w_0} \geq \lambda > 1$.

By Lemma 2.1, one has that the invariant set Λ is uniformly hyperbolic. The proof is completed. □

Therefore, it follows from Lemmas 2.2, 3.8, and 3.9 that Theorem 3.5 holds.

Based on the proof of Theorems 3.4 and 3.5, one has the following result:

Theorem 3.6. Suppose that there are $1 \leq i_0 < r_1$ and $1 \leq j_0 < r_2$ such that $0 = \alpha_{i_0} < \alpha_{i_0+1}$ and $0 < \xi_{j_0} < \xi_{j_0+1}$, and $m_{i_0} = m_{i_0+1} = 1$. Suppose that

(1). $a_2 < 0$ and $a_1 p'(0) > 0$;

(2). $\max\{0, b_3\alpha_{i_0+1}, b_1\alpha_{i_0+1}, b_1\alpha_{i_0+1} + b_3\alpha_{i_0+1}\} < \xi_{j_0}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) > 0$.

Then, for fixed a_2, b_1, and b_3, and sufficiently large $|a_1|$ and $|b_2|$, there exists a Smale horseshoe for the map (3.17), especially, there is a uniformly hyperbolic invariant set Λ on which F is topologically conjugate to the two-sided fullshift on four symbols. Therefore, they are Li-Yorke chaotic as well as Devaney chaotic.

Theorem 3.7. For system (3.17), suppose that $p(x) = \prod_{i=1}^{m}(x-\alpha_i)$ and $q(y) = \prod_{j=1}^{n}(y-\xi_j)$, where $\alpha_1 < \alpha_2 < \cdots < \alpha_m$ and $\xi_1 < \xi_2 < \cdots < \xi_n$ are real numbers, $m \geq 2$, and $n \geq 2$. Then, for fixed a_2, b_1, and b_3, and sufficiently large $|a_1|$ and $|b_2|$, there exist a Smale horseshoe for the map (3.17) and a uniformly hyperbolic invariant set Λ on which F is topologically conjugate to the two-sided fullshift on mn symbols. Therefore, they are chaotic in the sense of both Li-Yorke and Devaney.

Proof. Fix a constant $\lambda > 1$. It follows from the properties of $p(x)$ and $q(y)$ that there exist positive constants η_i, $1 \leq i \leq m$, and τ_j, $1 \leq j \leq n$, such that, the sets $V_i = [\alpha_i - \eta_i, \alpha_i + \eta_i]$, $1 \leq i \leq m$, are pairwise disjoint, and $|p'(x)| > 0$ for all $x \in V_i$, $1 \leq i \leq m$; the sets $W_j = [\xi_j - \tau_j, \xi_j + \tau_j]$, $1 \leq j \leq n$, are pairwise disjoint, and $|q'(y)| > 0$ for all $y \in W_j$, $1 \leq j \leq n$.

Denote

$$U := [\alpha_1 - \eta_1, \alpha_m + \eta_m] \times [\xi_1 - \tau_1, \xi_n + \tau_n] \times [\alpha_1 - \eta_1, \alpha_m + \eta_m],$$

and

$$\Lambda := \bigcap_{-\infty}^{+\infty} F^j(U).$$

Set

$$M_0 := \inf_{x \in \cup_{i=1}^{m} V_i} |p'(x)|, \ M_0' := \inf_{y \in \cup_{j=1}^{n} W_j} |q'(y)|;$$

$$M_1 := \min_{1\le i\le m} |p(\alpha_i \pm \eta_i)|,\ \ M_1' := \min_{1\le j\le n} |q(\xi_j \pm \tau_j)|;$$

$$N_0 := \frac{\max\{\lambda+|a_2|, 1+|a_2b_3|\lambda+|a_2|(1+|b_1|)\}}{M_0},\ \ N_0' := \frac{\lambda+|b_1|+|b_3|}{M_0'};$$

$$N_1 := \frac{|a_2|\max\{|\xi_1-\tau_1|, |\xi_n-\tau_n|\}+\max\{|\alpha_1-\eta_1|, |\alpha_m-\eta_m|\}}{M_1},$$

$$N_1' := \frac{\max\{|\xi_1-\tau_1|, |\xi_n-\tau_n|\}+(|b_1|+|b_3|)\max\{|\alpha_1-\eta_1|, |\alpha_m-\eta_m|\}}{M_1'}.$$

If $|a_1| > N_0$ and $|b_2| > N_0'$, by applying similar approaches in the proof of Lemma 3.7, one has that the invariant set Λ is uniformly hyperbolic, where the constants N_0 and N_0' have similar effects with the constants specified in (3.22).

If $|a_1| > N_1$ and $|b_2| > N_1'$, it follows from simple computation and the fact that $p(x)$ and $q(y)$ have simple real roots, one has that $F(U) \cap U$ has mn connected components.

Hence, if $|a_1| > \max\{N_0, N_1\}$ and $|b_2| > \max\{N_0', N_1'\}$, then there exist a Smale horseshoe and a hyperbolic invariant set on which F is topologically conjugate to the two-sided fullshift on mn symbols. This, together with Lemma 2.2, yields that F is Li-Yorke chaotic as well as Devaney chaotic. This completes the proof.

□

Remark 3.3. Similar results could be obtained for different parameters. For example, we could assume that $b_3 = c_1 = c_3 = 0$ and $a_3b_1 \neq 0$.

3.3 The expanding maps

In this subsection, we consider the following type of maps: $F : (x, y, z) \in \mathbb{R}^3 \to (f_1, f_2, f_3) \in \mathbb{R}^3$:

$$\begin{cases} f_1(x,y,z) = a_1p(x) + a_2y + a_3z \\ f_2(x,y,z) = b_1x + b_2q(y) + b_3z \\ f_3(x,y,z) = c_1x + c_2y + c_3r(z) \end{cases}, \tag{3.37}$$

where $a_1b_2c_3 \neq 0$.

The derivative of (3.37) is

$$DF = \begin{pmatrix} a_1p'(x) & a_2 & a_3 \\ b_1 & b_2q'(y) & b_3 \\ c_1 & c_2 & c_3r'(z) \end{pmatrix}. \tag{3.38}$$

Generally, the map (3.37) is not invertible.

An example is given as follows:

$$\begin{cases} f_1(x,y,z) = a_1x(1-x) + a_2y + a_3z \\ f_2(x,y,z) = b_1x + b_2y(1-y) + b_3z \\ f_3(x,y,z) = c_1x + c_2y + c_3z(1-z) \end{cases}.$$

This example can also be thought of as the generalization of the Hénon map in three-dimensional spaces.

Theorem 3.8. Suppose that there are $1 \le i_0 < r_1$, $1 \le j_0 < r_2$, and $1 \le k_0 < r_3$ such that $0 = \alpha_{i_0} < \alpha_{i_0+1}$, $0 = \xi_{j_0} < \xi_{j_0+1}$, and $0 = \kappa_{k_0} < \kappa_{k_0+1}$, and $m_{i_0} = m_{i_0+1} = n_{j_0} = n_{j_0+1} = l_{k_0} = l_{k_0+1} = 1$, that is, there are two distinct non-negative zeros of $p(x)$, $q(y)$, and $r(z)$. Suppose that

(1). $a_2 \le 0,\ a_3 \le 0,$ and $a_1p'(0) > 0$;

(2). $b_1 \le 0,\ b_3 \le 0,$ and $b_2q'(0) > 0$;

(3). $c_1 \le 0,\ c_2 \le 0,$ and $c_3r'(0) > 0$.

For fixed a_2, a_3, b_1, b_3, c_1, and c_2, if $|a_1|$, $|b_2|$, and $|c_3|$ are sufficiently large, then there exists a forward invariant set of the map (3.37) on which the map is topologically semi-conjugate to the one-sided fullshift on eight symbols and it is chaotic in the sense of Li-Yorke.

Proof. Denote

$$U := [0, \alpha_{i_0+1}] \times [0, \xi_{j_0+1}] \times [0, \kappa_{k_0+1}].$$

Fix $\lambda > 1$, a_2, a_3, b_1, b_3, c_1, and c_2. It follows from Lemma 3.1 that one could assume that $|a_1|$, $|b_2|$, and $|c_3|$ are sufficiently large such that for $w = (x,y,z) \in U \cap F^{-1}(U)$, one has that $x \in [0,\delta_1] \cup [\delta_1', \alpha_{i_0+1}]$, $y \in [0,\delta_2] \cup [\delta_2', \xi_{j_0+1}]$, and $z \in [0,\delta_3] \cup [\delta_3', \kappa_{k_0+1}]$.

Set

$$M_0 := \min\left\{ \min_{x\in[0,\delta_1]\cup[\delta_1',\alpha_{i_0+1}]} |p'(x)|,\ \min_{y\in[0,\delta_2]\cup[\delta_2',\xi_{j_0+1}]} |q'(y)|,\ \min_{z\in[0,\delta_3]\cup[\delta_3',\kappa_{k_0+1}]} |r'(z)| \right\}.$$

Suppose that

$$|a_1| \ge \frac{\lambda + |a_2| + |a_3|}{M_0},\ |b_2| \ge \frac{\lambda + |b_1| + |b_3|}{M_0},\ |c_3| \ge \frac{\lambda + |c_1| + |c_2|}{M_0}. \tag{3.39}$$

For $v = (v_{01}, v_{02}, v_{03})^T \in \mathbb{R}^3$, set $|v| = \max_{1\le i\le 3} |v_{0i}|$. By (3.38),

$$\begin{cases} v_{11} = a_1p'(x_0)v_{01} + a_2v_{02} + a_3v_{03} \\ v_{12} = b_1v_{01} + b_2q'(y_0)v_{02} + b_3v_{03} \\ v_{13} = c_1v_{01} + c_2v_{02} + c_3r'(z_0)v_{03} \end{cases}.$$

Without loss of generality, suppose that $|v| = |v_{01}|$. Hence, by (3.39), one has

$$|v_1| = \max_{1\le i\le 3} |v_{1i}| \ge |v_{11}| \ge |a_1p'(x_0)||v| - |a_2||v| - |a_3||v| \ge \lambda|v|,$$

which implies that $\|DF\| \ge \lambda$. So, one has that the map is expansion on $U \cap F^{-1}(U)$.

For fixed $a_2, a_3, b_1, b_3, c_1, c_2$, if $|a_1|$, $|b_2|$, and $|c_3|$ are sufficiently large, then there exist eight different closed subsets $V_1, ..., V_8$ of U such that $V_i \cap V_j = \emptyset$, $1 \le i \ne j \le 8$, and $F(V_i) = U$. This is a coupled-expanding map [54, 55]. By Theorem 3.1 in [55], there exists a forward invariant set on which the map is topologically semi-conjugate to the one-sided fullshift on eight symbols, and it is chaotic in the sense of Li-Yorke. This completes the proof. □

Theorem 3.9. Suppose that there are $1 \le i_0 < r_1$, $1 \le j_0 < r_2$, and $1 \le k_0 < r_3$ such that $0 < \alpha_{i_0} < \alpha_{i_0+1}$, $0 < \xi_{j_0} < \xi_{j_0+1}$, and $0 < \kappa_{k_0} < \kappa_{k_0+1}$, that is, there are two distinct positive zeros of $p(x)$, $q(y)$, and $r(z)$. Set

$$M_1 := \max\{a_2\xi_{j_0} + a_3\kappa_{k_0}, a_2\xi_{j_0+1} + a_3\kappa_{k_0}, a_2\xi_{j_0} + a_3\kappa_{k_0+1}, a_2\xi_{j_0+1} + a_3\kappa_{k_0+1}\},$$

$$M_2 := \min\{a_2\xi_{j_0} + a_3\kappa_{k_0}, a_2\xi_{j_0+1} + a_3\kappa_{k_0}, a_2\xi_{j_0} + a_3\kappa_{k_0+1}, a_2\xi_{j_0+1} + a_3\kappa_{k_0+1}\},$$

$$N_1 := \max\{b_1\alpha_{i_0} + b_3\kappa_{k_0}, b_1\alpha_{i_0+1} + b_3\kappa_{k_0}, b_1\alpha_{i_0} + b_3\kappa_{k_0+1}, b_1\alpha_{i_0+1} + b_3\kappa_{k_0+1}\},$$

$$N_2 := \min\{b_1\alpha_{i_0} + b_3\kappa_{k_0}, b_1\alpha_{i_0+1} + b_3\kappa_{k_0}, b_1\alpha_{i_0} + b_3\kappa_{k_0+1}, b_1\alpha_{i_0+1} + b_3\kappa_{k_0+1}\},$$

$$Q_1 := \max\{c_1\alpha_{i_0} + c_2\xi_{j_0}, c_1\alpha_{i_0+1} + c_2\xi_{j_0}, c_1\alpha_{i_0} + c_2\xi_{j_0+1}, c_1\alpha_{i_0+1} + c_2\xi_{j_0+1}\},$$

$$Q_2 := \min\{c_1\alpha_{i_0} + c_2\xi_{j_0}, c_1\alpha_{i_0+1} + c_2\xi_{j_0}, c_1\alpha_{i_0} + c_2\xi_{j_0+1}, c_1\alpha_{i_0+1} + c_2\xi_{j_0+1}\}.$$

Consider the following different situations:

(i). $M_1 < \alpha_{i_0}$ and $a_1p^{(m_{i_0})}(\alpha_{i_0}) > 0$, $N_1 < \xi_{j_0}$ and $b_2q^{(n_{j_0})}(\xi_{j_0}) > 0$, and $Q_1 < \kappa_{k_0}$ and $c_3r^{(l_{k_0})}(\kappa_{k_0}) > 0$;

(ii). $M_1 < \alpha_{i_0}$ and $a_1p^{(m_{i_0})}(\alpha_{i_0}) > 0$, $N_1 < \xi_{j_0}$ and $b_2q^{(n_{j_0})}(\xi_{j_0}) > 0$, and $Q_2 > \kappa_{k_0+1}$ and $c_3r^{(l_{k_0})}(\kappa_{k_0}) < 0$;

(iii). $M_1 < \alpha_{i_0}$ and $a_1 p^{(m_{i_0})}(\alpha_{i_0}) > 0$, $N_2 > \xi_{j_0+1}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) < 0$, and $Q_1 < \kappa_{k_0}$ and $c_3 r^{(l_{k_0})}(\kappa_{k_0}) > 0$;

(iv). $M_1 < \alpha_{i_0}$ and $a_1 p^{(m_{i_0})}(\alpha_{i_0}) > 0$, $N_2 > \xi_{j_0+1}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) < 0$, and $Q_2 > \kappa_{k_0+1}$ and $c_3 r^{(l_{k_0})}(\kappa_{k_0}) < 0$;

(v). $M_2 > \alpha_{i_0+1}$ and $a_1 p^{(m_{i_0})}(\alpha_{i_0}) < 0$, $N_1 < \xi_{j_0}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) > 0$, and $Q_1 < \kappa_{k_0}$ and $c_3 r^{(l_{k_0})}(\kappa_{k_0}) > 0$;

(vi). $M_2 > \alpha_{i_0+1}$ and $a_1 p^{(m_{i_0})}(\alpha_{i_0}) < 0$, $N_1 < \xi_{j_0}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) > 0$, and $Q_2 > \kappa_{k_0+1}$ and $c_3 r^{(l_{k_0})}(\kappa_{k_0}) < 0$;

(vii). $M_2 > \alpha_{i_0+1}$ and $a_1 p^{(m_{i_0})}(\alpha_{i_0}) < 0$, $N_2 > \xi_{j_0+1}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) < 0$, and $Q_1 < \kappa_{k_0}$ and $c_3 r^{(l_{k_0})}(\kappa_{k_0}) > 0$;

(viii). $M_2 > \alpha_{i_0+1}$ and $a_1 p^{(m_{i_0})}(\alpha_{i_0}) < 0$, $N_2 > \xi_{j_0+1}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) < 0$, and $Q_2 > \kappa_{k_0+1}$ and $c_3 r^{(l_{k_0})}(\kappa_{k_0}) < 0$.

For fixed a_2, a_3, b_1, b_3, c_1, c_2, if $|a_1|$, $|b_2|$, and $|c_3|$ are sufficiently large, then there exists a forward invariant set of the map (3.37) on which the map is topologically semi-conjugate to the one-sided fullshift on eight symbols and it is chaotic in the sense of Li-Yorke.

Remark 3.4. The eight different cases in Theorem 3.9 are the generalization of the assumptions (3.10) in Theorem 3.2 and the four situations in the assumptions of Theorem 3.5.

Set

$$U := [\alpha_{i_0}, \alpha_{i_0+1}] \times [\xi_{j_0}, \xi_{j_0+1}] \times [\kappa_{k_0}, \kappa_{k_0+1}].$$

Fix a positive constant $\lambda > 1$.

Proof. Fix a_2, a_3, b_1, b_3, c_1, and c_2. By Lemma 3.1, one could assume that $|a_1|$, $|b_2|$, and $|c_3|$ are sufficiently large such that for $w = (x, y, z) \in U \cap F^{-1}(U)$, one has that $x \in (\alpha_{i_0}, \delta_1] \cup [\delta'_1, \alpha_{i_0+1})$, $y \in (\xi_{j_0}, \delta_2] \cup [\delta'_2, \xi_{j_0+1})$, and $z \in (\kappa_{k_0}, \delta_3] \cup [\delta'_3, \kappa_{k_0+1})$.

In the following discussions, we only study Case (i), other cases could be treated by applying similar methods.

By (3.1)–(3.3), if $|a_1|$, $|b_2|$, and $|c_3|$ are sufficiently large, then for any $w = (x, y, z) \in U \cap F^{-1}(U)$,

$$\frac{|a_1 p'(x)|}{|a_1 p(x)|} \geq \frac{\lambda + |a_2| + |a_3|}{a_{i_0} - M_1}, \frac{|b_2 q'(y)|}{|b_2 q(y)|} \geq \frac{\lambda + |b_1| + |b_3|}{\xi_{j_0} - N_1}, \frac{|c_3 r'(z)|}{|c_3 r(z)|} \geq \frac{\lambda + |c_1| + |c_2|}{\kappa_{k_0} - Q_1}. \tag{3.40}$$

For any point $w_0 = (x_0, y_0, z_0) \in U \cap F^{-1}(U)$, set $w_1 = F(w_0) = (x_1, y_1, z_1)$. By (3.37),

$$\left\{\begin{array}{l} x_1 = a_1 p(x_0) + a_2 y_0 + a_3 z_0 \\ y_1 = b_1 x_0 + b_2 q(y_0) + b_3 z_0 \\ z_1 = c_1 x_0 + c_2 y_0 + c_3 r(z_0) \end{array}\right. .$$

So, one has that $a_1 p(x_0) = x_1 - a_2 y_0 - a_3 z_0$, $b_2 q(y_0) = y_1 - b_1 x_0 - b_3 z_0$, and $c_3 r(z_0) = z_1 - c_1 x_0 - c_2 y_0$. This yields that for any $w = (x, y, z) \in U \cap F^{-1}(U)$,

$$a_1 p(x) \geq \alpha_{i_0} - M_1,\ b_2 q(y) \geq \xi_{j_0} - N_1,\ c_3 r(z) \geq \kappa_{k_0} - Q_1. \tag{3.41}$$

By (3.40) and (3.41), one has that for any $w = (x, y, z) \in U \cap F^{-1}(U)$,

$$|a_1 p'(x)| \geq \lambda + |a_2| + |a_3|,\ |b_2 q'(y)| \geq \lambda + |b_1| + |b_3|,\ |c_3 r'(z)| \geq \lambda + |c_1| + |c_2|. \tag{3.42}$$

Next, it is to show that the map in Case (i) is expansion in distance.

For $v = (v_{01}, v_{02}, v_{03})^T \in \mathbb{R}^3$, set $|v| = \max_{1 \leq i \leq 3} |v_{0i}|$. By (3.38),

$$\left\{\begin{array}{l} v_{11} = a_1 p'(x_0) v_{01} + a_2 v_{02} + a_3 v_{03} \\ v_{12} = b_1 v_{01} + b_2 q'(y_0) v_{02} + b_3 v_{03} \\ v_{13} = c_1 v_{01} + c_2 v_{02} + c_3 r'(z_0) v_{03} \end{array}\right. .$$

Without loss of generality, suppose that $|v| = |v_{01}|$. Hence, From (3.42), it follows that

$$|v_1| = \max_{1 \leq i \leq 3} |v_{1i}| \geq |v_{11}| \geq |a_1 p'(x_0)||v| - |a_2||v| - |a_3||v| \geq \lambda |v|,$$

which implies that $\|DF\| \geq \lambda$. So, one has that the map is expansion on $U \cap F^{-1}(U)$.

Hence, there exist eight disjoint closed subsets $V_1, ..., V_8$ of U such that $F(V_i) = U$, $1 \leq i \leq 8$. The map F on $\cup_{1 \leq i \leq 8} V_i$ is a coupled-expanding map [54, 55]. It follows from Theorem 3.1 in [55] that there is a forward invariant set on which the map is topologically semi-conjugate to the one-sided fullshift on eight symbols, and it is chaotic in the sense of Li-Yorke. The proof is completed. □

By the proof of Theorems 3.8 and 3.9, one has the following results:

Theorem 3.10. Suppose that there are $1 \leq i_0 < r_1$, $1 \leq j_0 < r_2$, and $1 \leq k_0 < r_3$ such that $0 = \alpha_{i_0} < \alpha_{i_0+1}$, $0 < \xi_{j_0} < \xi_{j_0+1}$, and $0 < \kappa_{k_0} < \kappa_{k_0+1}$, and $m_{i_0} = m_{i_0+1} = 1$. Set

$$N_1 := \max\{b_3\kappa_{k_0}, b_1\alpha_{i_0+1} + b_3\kappa_{k_0}, b_3\kappa_{k_0+1}, b_1\alpha_{i_0+1} + b_3\kappa_{k_0+1}\},$$

$$N_2 := \min\{b_3\kappa_{k_0}, b_1\alpha_{i_0+1} + b_3\kappa_{k_0}, b_3\kappa_{k_0+1}, b_1\alpha_{i_0+1} + b_3\kappa_{k_0+1}\},$$

$$Q_1 := \max\{c_2\xi_{j_0}, c_1\alpha_{i_0+1} + c_2\xi_{j_0}, c_2\xi_{j_0+1}, c_1\alpha_{i_0+1} + c_2\xi_{j_0+1}\},$$

$$Q_2 := \min\{c_2\xi_{j_0}, c_1\alpha_{i_0+1} + c_2\xi_{j_0}, c_2\xi_{j_0+1}, c_1\alpha_{i_0+1} + c_2\xi_{j_0+1}\}.$$

Consider the following four different cases:

(i). $a_2 \leq 0$, $a_3 \leq 0$, and $a_1 p'(0) > 0$, $N_1 < \xi_{j_0}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) > 0$, and $Q_1 < \kappa_{k_0}$ and $c_3 r^{(l_{k_0})}(\kappa_{k_0}) > 0$;

(ii). $a_2 \leq 0$, $a_3 \leq 0$, and $a_1 p'(0) > 0$, $N_1 < \xi_{j_0}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) > 0$, and $Q_2 > \kappa_{k_0+1}$ and $c_3 r^{(l_{k_0})}(\kappa_{k_0}) < 0$;

(iii). $a_2 \leq 0$, $a_3 \leq 0$, and $a_1 p'(0) > 0$, $N_2 > \xi_{j_0+1}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) < 0$, and $Q_1 < \kappa_{k_0}$ and $c_3 r^{(l_{k_0})}(\kappa_{k_0}) > 0$;

(iv). $a_2 \leq 0$, $a_3 \leq 0$, and $a_1 p'(0) > 0$, $N_2 > \xi_{j_0+1}$ and $b_2 q^{(n_{j_0})}(\xi_{j_0}) < 0$, and $Q_2 > \kappa_{k_0+1}$ and $c_3 r^{(l_{k_0})}(\kappa_{k_0}) < 0$.

For fixed a_2, a_3, b_1, b_3, c_1, and c_2, if $|a_1|$, $|b_2|$, and $|c_3|$ are sufficiently large, then there exists a forward invariant set of the map (3.37) on which the map is topologically semi-conjugate to the one-sided fullshift on eight symbols and it is chaotic in the sense of Li-Yorke.

Theorem 3.11. Suppose that there are $1 \leq i_0 < r_1$, $1 \leq j_0 < r_2$, and $1 \leq k_0 < r_3$ such that $0 = \alpha_{i_0} < \alpha_{i_0+1}$, $0 = \xi_{j_0} < \xi_{j_0+1}$, and $0 < \kappa_{k_0} < \kappa_{k_0+1}$, and $m_{i_0} = m_{i_0+1} = n_{j_0} = n_{j_0+1} = 1$. Suppose that

(1). $a_2 \leq 0$, $a_3 \leq 0$, and $a_1 p'(0) > 0$;

(2). $b_1 \leq 0$, $b_3 \leq 0$, and $b_2 q'(0) > 0$;

(3). $\max\{0, c_1\alpha_{i_0+1}, c_2\xi_{j_0+1}, c_1\alpha_{i_0+1} + c_2\xi_{j_0+1}\} < \kappa_{k_0}$ and $c_3 r^{(l_{k_0})}(\kappa_{k_0}) > 0$.

For fixed a_2, a_3, b_1, b_3, c_1, c_2, if $|a_1|$, $|b_2|$, and $|c_3|$ are sufficiently large, then there exists a forward invariant set of the map (3.37) on which the map is topologically semi-conjugate to the one-sided fullshift on eight symbols and it is chaotic in the sense of Li-Yorke.

Remark 3.5. Similar results could be obtained if the polynomials $p(x)$, $q(y)$, and $r(z)$ have two different non-positive roots, respectively.

4 Smale horseshoe in high-dimensional polynomial maps

In this section, we study the existence of Smale horseshoe and uniformly hyperbolic invariant sets of the maps (1.4), we generalize the well-known results obtained in [17].

The derivative of the map (1.4) is

$$DF = \mathrm{diag}[-2x_1, -2x_2, ..., -2x_d, \underbrace{0, ..., 0}_{n-d}] + \begin{pmatrix} 0 & a_{12} & \cdots & a_{1n} \\ a_{21} & 0 & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{n1} & a_{n2} & \cdots & 0 \end{pmatrix}. \tag{4.1}$$

Now, it is to study the inverse of the map F. We consider the following three types of maps with the inverse expressions.

Case (1). Suppose that $n \geq 2d$, and (1.4) can be written as follows:

$$\begin{pmatrix} f_1(x_1, ..., x_n) \\ \vdots \\ f_d(x_1, ..., x_n) \\ f_{d+1}(x_1, ..., x_n) \\ \vdots \\ f_n(x_1, ..., x_n) \end{pmatrix} = \begin{pmatrix} a_{11} - x_1^2 \\ \vdots \\ a_{dd} - x_d^2 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \begin{array}{c c} & \begin{array}{ccc} d & n-2d & d \end{array} \\ \begin{array}{c} d \\ n-d \end{array} & \begin{pmatrix} A & B & C \\ D & E & 0 \end{pmatrix} \end{array} \begin{pmatrix} x_1 \\ \vdots \\ x_d \\ x_{d+1} \\ \vdots \\ x_n \end{pmatrix}, \tag{4.2}$$

where C and $G = (D, E)$ are invertible matrices.

By direct calculation, one has that the inverse map $H : \mathbb{R}^n \to \mathbb{R}^n$ is

$$\begin{pmatrix} h_1(x_1, ..., x_n) \\ \vdots \\ h_{n-d}(x_1, ..., x_n) \end{pmatrix} = G^{-1} \begin{pmatrix} x_{d+1} \\ \vdots \\ x_n \end{pmatrix}, \tag{4.3}$$

$$\begin{pmatrix} h_{n-d+1}(x_1, ..., x_n) \\ \vdots \\ h_n(x_1, ..., x_n) \end{pmatrix} = C^{-1} \begin{pmatrix} x_1 \\ \vdots \\ x_d \end{pmatrix} - C^{-1} L \begin{pmatrix} h_1 \\ \vdots \\ h_d \end{pmatrix} - C^{-1} \begin{pmatrix} a_{11} - h_1^2 \\ a_{22} - h_2^2 \\ \vdots \\ a_{dd} - h_d^2 \end{pmatrix}, \tag{4.4}$$

where $L = (A, B)$ is a $d \times (n-d)$ matrix. In the above expressions, we should use (4.3) to substitute $h_1, ..., h_d$ in the right hand side of (4.4).

Suppose that

$$G^{-1} = (D, E)^{-1} = \begin{array}{c} \\ d \\ n-2d \end{array} \begin{array}{c} n-d \\ \begin{pmatrix} P \\ Q \end{pmatrix} \end{array}. \tag{4.5}$$

The Jacobian of the inverse map is

$$DH = \begin{array}{c} \\ n-d \\ d \end{array} \begin{array}{c} \begin{array}{cc} d & \qquad n-d \end{array} \\ \begin{pmatrix} 0 & G^{-1} \\ C^{-1} & -C^{-1}LP + S \end{pmatrix} \end{array}, \tag{4.6}$$

where

$$S = C^{-1}(\mathrm{diag}[2h_1, ..., 2h_d])P.$$

Case (2). Suppose that $\frac{3}{2}d \le n < 2d$, and (1.4) can be written as follows:

$$\begin{pmatrix} f_1(x_1, ..., x_n) \\ \vdots \\ f_d(x_1, ..., x_n) \\ f_{d+1}(x_1, ..., x_n) \\ \vdots \\ f_n(x_1, ..., x_n) \end{pmatrix} = \begin{pmatrix} a_{11} - x_1^2 \\ \vdots \\ a_{dd} - x_d^2 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \begin{array}{c} \\ 2d-n \\ n-d \\ n-d \end{array} \begin{array}{c} \begin{array}{ccc} n-d & 2d-n & n-d \end{array} \\ \begin{pmatrix} A & B & 0 \\ C & D & E \\ G & 0 & 0 \end{pmatrix} \end{array} \begin{pmatrix} x_1 \\ \vdots \\ x_d \\ x_{d+1} \\ \vdots \\ x_n \end{pmatrix}, \tag{4.7}$$

where B, E, and G are invertible matrices.

The inverse map $H : \mathbb{R}^n \to \mathbb{R}^n$ is as follows:

$$\begin{pmatrix} h_1 \\ \vdots \\ h_{n-d} \end{pmatrix} = G^{-1} \begin{pmatrix} x_{d+1} \\ \vdots \\ x_n \end{pmatrix},$$

$$\begin{pmatrix} h_{n-d+1} \\ \vdots \\ h_d \end{pmatrix} = B^{-1} \left(\begin{pmatrix} x_1 \\ \vdots \\ x_{2d-n} \end{pmatrix} - \begin{pmatrix} a_{11} - h_1^2 \\ \vdots \\ a_{2d-n,2d-n} - h_{2d-n}^2 \end{pmatrix} - AG^{-1} \begin{pmatrix} x_{d+1} \\ \vdots \\ x_n \end{pmatrix} \right),$$

$$\begin{pmatrix} h_{d+1} \\ \vdots \\ h_n \end{pmatrix} = E^{-1}\left(\begin{pmatrix} x_{2d-n+1} \\ \vdots \\ x_d \end{pmatrix} - \begin{pmatrix} a_{2d-n+1,2d-n+1} - h_{2d-n+1}^2 \\ \vdots \\ a_{dd} - h_d^2 \end{pmatrix} - C \begin{pmatrix} h_1 \\ \vdots \\ h_{2d-n} \end{pmatrix}\right) - E^{-1}D \begin{pmatrix} h_{2d-n+1} \\ \vdots \\ h_d \end{pmatrix}.$$

It is evident that the degree of the inverse map is four.

Suppose that

$$G^{-1} = \begin{array}{c} \\ 2d-n \\ 2n-3d \end{array}\!\!\begin{array}{c} n-d \\ \left(\begin{array}{c} K \\ L \end{array}\right) \end{array}, \quad E^{-1} = \begin{array}{c} \\ n-d \end{array}\!\!\begin{array}{c} \begin{array}{cc} 2n-3d & 2d-n \end{array} \\ \left(\begin{array}{cc} M & N \end{array}\right) \end{array},$$

and

$$E^{-1}D = \begin{array}{c} \\ n-d \end{array}\!\!\begin{array}{c} \begin{array}{cc} 2n-3d & 2d-n \end{array} \\ \left(\begin{array}{cc} P & Q \end{array}\right) \end{array}.$$

The derivative of the map is

$$\begin{array}{c|ccc|} & 2d-n & n-d & n-d \\ n-d & 0 & 0 & G^{-1} \\ 2d-n & B^{-1} & 0 & S \\ n-d & V & E^{-1} & W \end{array}, \tag{4.8}$$

where

$$S = B^{-1}(\mathrm{diag}[2h_1, ..., 2h_{2d-n}])K - B^{-1}AG^{-1},$$
$$V = -QB^{-1} + N(\mathrm{diag}[2h_{n-d+1}, ..., 2h_d])B^{-1},$$
$$W = E^{-1}(\mathrm{diag}[2h_{2d-n+1}, ..., 2h_d]) \begin{pmatrix} L \\ S \end{pmatrix} - E^{-1}CK - E^{-1}D \begin{pmatrix} L \\ S \end{pmatrix}.$$

Case (3). It is to investigate the following type of map, where $n < 2d$ and the degree of the inverse map is equal to 2^{2d-n}.

$$\begin{pmatrix} f_1(x_1, ..., x_n) \\ \vdots \\ f_d(x_1, ..., x_n) \\ f_{d+1}(x_1, ..., x_n) \\ \vdots \\ f_n(x_1, ..., x_n) \end{pmatrix} = \begin{pmatrix} a_{11} - x_1^2 \\ \vdots \\ a_{dd} - x_d^2 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \begin{array}{c} \\ n-d-1 \\ 2d-n+1 \\ n-d \end{array}\!\!\begin{array}{c} \begin{array}{ccc} n-d & 2d-n+1 & n-d-1 \end{array} \\ \left(\begin{array}{ccc} A & B & C \\ D & E & 0 \\ G & 0 & 0 \end{array}\right) \end{array} \begin{pmatrix} x_1 \\ \vdots \\ x_d \\ x_{d+1} \\ \vdots \\ x_n \end{pmatrix}, \tag{4.9}$$

where C and G are invertible. In the matrix $\begin{pmatrix} B \\ E \end{pmatrix}$, $\prod_{i=0}^{2d-n} a_{n-d+i,n-d+i+1} \neq 0$, all the other coefficients are zero. Suppose that

$$D = \begin{pmatrix} s_1 \\ \vdots \\ s_{2d-n+1} \end{pmatrix}, \tag{4.10}$$

where s_i is a $1 \times (n-d)$ matrix, $1 \leq i \leq 2d-n+1$.

The inverse map $H : \mathbb{R}^n \to \mathbb{R}^n$ is

$$\begin{pmatrix} h_1 \\ \vdots \\ h_{n-d} \end{pmatrix} = G^{-1} \begin{pmatrix} x_{d+1} \\ \vdots \\ x_n \end{pmatrix}, \tag{4.11}$$

$$\begin{pmatrix} h_{d+2} \\ \vdots \\ h_n \end{pmatrix} = C^{-1} \left(\left(\begin{pmatrix} x_1 \\ \vdots \\ x_{n-d-1} \end{pmatrix} - \begin{pmatrix} a_{11} - h_1^2 \\ \vdots \\ a_{n-d-1,n-d-1} - h_{n-d-1}^2 \end{pmatrix} \right) - AG^{-1} \begin{pmatrix} x_{d+1} \\ \vdots \\ x_n \end{pmatrix} \right);$$

the map h_{n-d+i} is given recurrently, $1 \leq i \leq 2d-n+1$,

$$h_{n-d+1} = \frac{1}{a_{n-d,n-d+1}} \left(x_{n-d} - (a_{n-d,n-d} - h_{n-d}^2) - s_1 \left(G^{-1} \begin{pmatrix} x_{d+1} \\ \vdots \\ x_n \end{pmatrix} \right) \right),$$

$$h_{n-d+i} = \frac{1}{a_{n-d+i-1,n-d+i}} \left(x_{n-d+i-1} - (a_{n-d+i-1,n-d+i-1} - h_{n-d+i-1}^2) - s_i G^{-1} \begin{pmatrix} x_{d+1} \\ \vdots \\ x_n \end{pmatrix} \right),$$
$$2 \leq i \leq 2d-n+1,$$

where s_i is specified in (4.10).

The degree of the inverse map is equal to 2^{2d-n}.

Suppose that

$$G^{-1} = \begin{array}{cc} & n-d \\ \begin{array}{c} n-d-1 \\ 1 \end{array} & \begin{pmatrix} K \\ L \end{pmatrix} \end{array}.$$

The derivative of the inverse map H is

$$\begin{array}{cc} & \begin{array}{ccc} n-d-1 & 2d-n+1 & n-d \end{array} \\ \begin{array}{c} n-d \\ 2d-n+1 \\ n-d-1 \end{array} & \begin{pmatrix} 0 & 0 & G^{-1} \\ 0 & S & -DG^{-1} \\ C^{-1} & 0 & V \end{pmatrix} \end{array} + \begin{array}{cc} & n \\ \begin{array}{c} n-d \\ 2d-n+1 \\ n-d-1 \end{array} & \begin{pmatrix} 0 \\ SW \begin{pmatrix} \frac{\partial h_{n-d}}{\partial x_1} & \cdots & \frac{\partial h_{n-d}}{\partial x_n} \\ \vdots & \vdots & \vdots \\ \frac{\partial h_d}{\partial x_1} & \cdots & \frac{\partial h_d}{\partial x_n} \end{pmatrix} \\ 0 \end{pmatrix} \end{array}, \tag{4.12}$$

where

$$S = \mathrm{diag}\left[\frac{1}{a_{n-d,n-d+1}}, ..., \frac{1}{a_{d,d+1}}\right],$$

$$V = C^{-1}(\mathrm{diag}[2h_1, ..., 2h_{n-d-1}])K - C^{-1}AG^{-1},$$

$$W = \mathrm{diag}[2h_{n-d}, ..., 2h_d],$$

the derivative of the inverse map is defined recurrently.

Set

$$R := \min_{1\le i\le d} \sqrt{a_{ii}},\ U := [-R, R]^n. \tag{4.13}$$

In the following discussions, fix a constant $\lambda > 1$.

Theorem 4.1. For the maps in Cases (1)–(3), for fixed a_{ij}, $1 \le i \ne j \le n$, and sufficiently large a_{11},...,a_{dd}, there exists a Smale horseshoe and a uniformly hyperbolic invariant set on which the map is topologically conjugate to the two-sided fullshift on 2^d symbols. Consequently, the map is chaotic in the sense of both Li-Yorke and Devaney.

First, it is to study the point $w_0 \in U \cap F(U)$, that is, $w_{-1} = F^{-1}(w_0) \in U$.

Lemma 4.1. In Case (1), for any fixed a_{ij} in (4.2), $1 \le i \ne j \le n$, there is a positive constant N_1, such that if $\min_{1\le i\le d}\{a_{ii}\} \ge N_1$, then for the map (4.2) and any $w_0 \in U \cap F(U)$, one has that $m'_{\mathcal{C},w_0} \ge \lambda > 1$.

Proof. For the point $w_0 \in U \cap F(U)$, set $w_{-1} = F^{-1}(w_0) = H(w_0) \in U$. By the invertibility of C, the expression of F^{-1}, and the definition of the region U, one has that

$$C_1 a_{ii} \le h_i^2 \le C_1' a_{ii},\ 1 \le i \le d, \tag{4.14}$$

where C_1 and C_1' are positive constants dependent on a_{ij}, $1 \le i \ne j \le n$, and $a_{11}, ..., a_{dd}$ are sufficiently large.

Consider the following unit cone:

$$K_1(\mathbb{R}^d, \mathbb{R}^{n-d}) = \Bigg\{ \mathbf{v} = \begin{pmatrix} v_0 \\ u_0 \end{pmatrix} : \ v_0 = (v_{0,1}, ..., v_{0,d})^T \in \mathbb{R}^d,$$
$$u_0 = (u_{0,1}, ..., u_{0,n-d})^T \in \mathbb{R}^{n-d}, \text{ and } |u_0| > |v_0| \Bigg\},$$

where $|v_0| = \max\{|v_{0,1}|, ..., |v_{0,d}|\}$, $|u_0| = \max\{|u_{0,1}|, ..., |u_{0,d}|\}$, and $|\mathbf{v}| = \max\{|v_0|, |u_0|\}$.

Suppose that $DH(\mathbf{v}) = \begin{pmatrix} v_{-1} \\ u_{-1} \end{pmatrix}$, where $v_{-1} = (v_{-1,1}, ..., v_{-1,d})^T \in \mathbb{R}^d$ and $u_{-1} = (u_{-1,1}, ..., u_{-1,n-d})^T \in \mathbb{R}^{n-d}$. It follows from (4.6) that

$$v_{-1} = G^{-1}u_0, \ u_{-1} = C^{-1}v_0 - C^{-1}LPu_0 + Su_0. \tag{4.15}$$

To show that $m'_{\mathcal{C},w_0} \geq \lambda$, it is sufficient to show that $|v_{-1}| < |u_{-1}|$ and $|u_{-1}| \geq \lambda |u_0|$. By (4.15), we only need to consider the following part:

$$Su_0 = S \begin{pmatrix} u_{0,1} \\ u_{0,2} \\ \vdots \\ u_{0,n-d} \end{pmatrix}.$$

Suppose that

$$P = \begin{pmatrix} s_1 \\ s_2 \\ \vdots \\ s_d \end{pmatrix},$$

where s_i is a $1 \times (n-d)$ matrix, $1 \leq i \leq d$. So, by direct calculations, one has that

$$S \begin{pmatrix} u_{0,1} \\ u_{0,2} \\ \vdots \\ u_{0,n-d} \end{pmatrix} = 2C^{-1} \begin{pmatrix} s_1 \begin{pmatrix} x_{d+1} \\ x_{d+2} \\ \vdots \\ x_n \end{pmatrix} s_1 \begin{pmatrix} u_{0,1} \\ u_{0,2} \\ \vdots \\ u_{0,n-d} \end{pmatrix} \\ \vdots \\ s_d \begin{pmatrix} x_{d+1} \\ x_{d+2} \\ \vdots \\ x_n \end{pmatrix} s_d \begin{pmatrix} u_{0,1} \\ u_{0,2} \\ \vdots \\ u_{0,n-d} \end{pmatrix} \end{pmatrix} = 2C^{-1} \begin{pmatrix} h_1 s_1 \begin{pmatrix} u_{0,1} \\ u_{0,2} \\ \vdots \\ u_{0,n-d} \end{pmatrix} \\ \vdots \\ h_d s_d \begin{pmatrix} u_{0,1} \\ u_{0,2} \\ \vdots \\ u_{0,n-d} \end{pmatrix} \end{pmatrix}. \tag{4.16}$$

This, together with (4.5) and (4.14), implies that if $a_{11}, ..., a_{dd}$ are sufficiently large, then $|u_{-1}| > |v_{-1}|$ and $|u_{-1}| \geq \lambda |u_0|$, yielding that $m'_{\mathcal{C},w_0} \geq \lambda > 1$. The proof is completed. □

Lemma 4.2. In Case (2), for any fixed a_{ij} in (4.7), $1 \leq i \neq j \leq n$, there is a positive constant N_1, such that if $\min_{1\leq i \leq d}\{a_{ii}\} \geq N_1$, then for the map (4.7) and any $w_0 \in U \cap F(U)$, one has that $m'_{\mathcal{C},w_0} \geq \lambda > 1$

Proof. It follows from the invertibility of B and E, the expression of the inverse of (4.7), and the definition of the region U, that

$$C_2 a_{ii} \leq h_i^2 \leq C'_2 a_{ii}, \; 1 \leq i \leq d, \tag{4.17}$$

where C_2 and C'_2 are positive constants dependent on a_{ij}, $1 \leq j \leq n$, and $j \neq i$, and $a_{11}, ..., a_{dd}$ are sufficiently large.

Consider the following unit cone:

$$\begin{aligned} K_1(\mathbb{R}^d, \mathbb{R}^{n-d}) = \Bigg\{ \mathbf{v} = \begin{pmatrix} v_0 \\ u_0 \end{pmatrix} : \; & v_0 = (v_{0,1}, ..., v_{0,d})^T \in \mathbb{R}^d, \\ & u_0 = (u_{0,1}, ..., u_{0,n-d})^T \in \mathbb{R}^{n-d}, \text{ and } |u_0| > |v_0| \Bigg\}, \end{aligned}$$

where $|v_0| = \max\{|v_{0,1}|, ..., |v_{0,d}|\}$, $|u_0| = \max\{|u_{0,1}|, ..., |u_{0,d}|\}$, and $|\mathbf{v}| = \max\{|v_0|, |u_0|\}$. Assume that $DH(\mathbf{v}) = \begin{pmatrix} v_{-1} \\ u_{-1} \end{pmatrix}$, where $v_{-1} = (v_{-1,1}, ..., v_{-1,d})^T \in \mathbb{R}^d$ and $u_{-1} = (u_{-1,1}, ..., u_{-1,n-d})^T \in \mathbb{R}^{n-d}$. Suppose that $v_0 = \begin{pmatrix} v_0^1 \\ v_0^2 \end{pmatrix}$ and $v_{-1} = \begin{pmatrix} v_{-1}^1 \\ v_{-1}^2 \end{pmatrix}$, where the dimension of v_0^1 and v_{-1}^1 is $2d - n$. By (4.8), one has that

$$\begin{pmatrix} v_{-1}^1 \\ v_{-1}^2 \\ u_{-1} \end{pmatrix} = \begin{pmatrix} G^{-1}u_0 \\ B^{-1}v_0^1 + Su_0 \\ Vv_0^1 + E^{-1}v_0^2 + Wu_0 \end{pmatrix}. \tag{4.18}$$

To show that $m'_{\mathcal{C},w_0} \geq \lambda$, it is sufficient to show that $|v_{-1}| < |u_{-1}|$ and $|u_{-1}| \geq \lambda |u_0|$. It follows from (4.8) and (4.18) that we only need to study Su_0, Vv_0^1, and Wu_0. So, we have to consider the following three terms:

$$\begin{aligned} & B^{-1}(\mathrm{diag}[2h_1, ..., 2h_{2d-n}])Ku_0, \; N(\mathrm{diag}[2h_{n-d+1}, ..., 2h_d])B^{-1}v_0^1, \\ & E^{-1}(\mathrm{diag}[2h_{2d-n+1}, ..., 2h_d]) \begin{pmatrix} L \\ B^{-1}(\mathrm{diag}[2h_1, ..., 2h_{2d-n}])K \end{pmatrix} u_0. \end{aligned}$$

Since $G^{-1} = \begin{pmatrix} K \\ L \end{pmatrix}$, where K is $(2d - n) \times (n - d)$, and L is $(2n - 3d) \times (n - d)$, one has that

$$\begin{aligned} & \left\| E^{-1}(\mathrm{diag}[2h_{2d-n+1}, ..., 2h_d]) \begin{pmatrix} L \\ B^{-1}(\mathrm{diag}[2h_1, ..., 2h_{2d-n}])K \end{pmatrix} \right\| \\ \gtrsim & \left\| B^{-1}(\mathrm{diag}[2h_1, ..., 2h_{2d-n}])K \right\|, \end{aligned}$$

if $a_{11}, ..., a_{dd}$ are sufficiently large. By direct calculation,

$$\begin{pmatrix} L \\ B^{-1}(\mathrm{diag}[2h_1, ..., 2h_{2d-n}])K \end{pmatrix} = \begin{pmatrix} E_{2n-3d} & 0 \\ 0 & B^{-1} \end{pmatrix} (\mathrm{diag}[\underbrace{1, ..., 1}_{2n-3d}, 2h_1, ..., 2h_{2d-n}]) \begin{pmatrix} L \\ K \end{pmatrix},$$

where $E_{2n-3d} = \mathrm{diag}[\underbrace{1, ..., 1}_{2n-3d}]$. It follows from the assumption $\frac{3}{2}d \leq n$, the fact that E and G are invertible, and (4.17), that

$$\|Wu_0\| \geq \|Vv_0^1\|,$$

if $a_{11}, ..., a_{dd}$ are sufficiently large.

Hence, $m'_{\mathcal{C},w_0} \geq \lambda > 1$ for sufficiently large $a_{11}, ..., a_{dd}$. This completes the proof. $\square$

Lemma 4.3. In Case (3), for any fixed a_{ij} in (4.9), $1 \leq i \neq j \leq n$, there is a positive constant N_1, such that if $\min_{1 \leq i \leq d}\{a_{ii}\} \geq N_1$, then for the map (4.9) and any $w_0 \in U \cap F(U)$, one has that $m'_{\mathcal{C},w_0} \geq \lambda > 1$.

Proof. By the invertibility of C and G, the expression of the inverse of (4.9), and the definition of the region U, one has that

$$C_3 a_{ii} \leq h_i^2 \leq C_3' a_{ii}, \ 1 \leq i \leq d, \tag{4.19}$$

where C_3 and C_3' are positive constants dependent on a_{ij}, $1 \leq i \neq j \leq n$, and $a_{11}, ..., a_{dd}$ are sufficiently large.

Consider the following unit cone:

$$K_1(\mathbb{R}^d, \mathbb{R}^{n-d}) = \Bigg\{ \mathbf{v} = \begin{pmatrix} v_0 \\ u_0 \end{pmatrix} : \ v_0 = (v_{0,1}, ..., v_{0,d})^T \in \mathbb{R}^d,$$
$$u_0 = (u_{0,1}, ..., u_{0,n-d})^T \in \mathbb{R}^{n-d}, \text{ and } |u_0| > |v_0| \Bigg\},$$

where $|v_0| = \max\{|v_{0,1}|, ..., |v_{0,d}|\}$, $|u_0| = \max\{|u_{0,1}|, ..., |u_{0,d}|\}$, and $|\mathbf{v}| = \max\{|v_0|, |u_0|\}$. Denote $DH(\mathbf{v}) = \begin{pmatrix} v_{-1} \\ u_{-1} \end{pmatrix}$, where $v_{-1} = (v_{-1,1}, ..., v_{-1,d})^T$ and $u_{-1} = (u_{-1,1}, ..., u_{-1,n-d})^T$. Suppose that $v_0 = \begin{pmatrix} v_0^1 \\ v_0^2 \end{pmatrix}$ with the dimension of v_0^1 equals to $n-d-1$, $u_{-1} = \begin{pmatrix} u_{-1}^1 \\ u_{-1}^2 \end{pmatrix}$ with the dimension of u_{-1}^1 equals to 1, $v_{-1} = \begin{pmatrix} v_{-1}^1 \\ v_{-1}^2 \end{pmatrix}$ with the dimension of v_{-1}^1 equals to $n-d$.

By (4.12), one has that

$$\left(\begin{array}{c} v_{-1}^1 \\ \hline v_{-1}^2 \\ u_{-1}^1 \\ \hline u_{-1}^2 \end{array}\right) = \left(\begin{array}{c} G^{-1}u_0 \\ \hline Sv_0^2 - DG^{-1}u_0 + SW \begin{pmatrix} \frac{\partial h_{n-d}}{\partial x_1} & \cdots & \frac{\partial h_{n-d}}{\partial x_n} \\ \vdots & \vdots & \vdots \\ \frac{\partial h_d}{\partial x_1} & \cdots & \frac{\partial h_d}{\partial x_n} \end{pmatrix} \mathbf{v} \\ \hline C^{-1}v_0^1 + Vu_0 \end{array}\right). \qquad (4.20)$$

To show that $m'_{\mathcal{C},w_0} \geq \lambda$, it is sufficient to show that $|v_{-1}| < |u_{-1}|$ and $|u_{-1}| \geq \lambda |u_0|$. By (4.12) and (4.20), one only need to investigate the expressions for v_{-1}^2, u_{-1}^1, and u_{-1}^2. By (4.11) and (4.12), if a_{11},...,a_{dd} are sufficiently large, then $|u_{-1}^1| \approx C\prod_{i=n-d}^{d}|h_i|$, where C is a constant dependent on $\mathbf{v}$ an the parameter a_{ij}, $i \neq j$. This, together with the expression for u_1^2 and (4.19), implies that $|v_{-1}| < |u_{-1}|$ and $|u_{-1}| \geq \lambda|u_0|$. This completes the whole proof. $\square$

Now, it is to study $U \cap F^{-1}(U)$.

Lemma 4.4. For any fixed a_{ij}, $1 \leq i \neq j \leq n$, there is a positive constant N_1, such that if $\min_{1\leq i\leq d}\{a_{ii}\} \geq N_1$, then for the map (1.4) and any $w_0 \in U \cap F^{-1}(U)$, one has that $m_{\mathcal{C},w_0} \geq \lambda > 1$.

Proof. Suppose that $w_0 = (x_{0,1}, ..., x_{0,n}) \in U$, $w_1 = F(w_0) = (x_{1,1}, ..., x_{1,n})$, and $w_{-1} = F^{-1}(w_0) = (x_{-1,1}, ..., x_{-1,n})$. Since $w_0 \in U \cap F^{-1}(U)$, one has that $w_1 = F(w_0) \in U$, that is,

$$-R \leq a_{ii} - x_i^2 + \sum_{1\leq j\leq n,\ j\neq i} a_{ij}x_j \leq R,\ 1 \leq i \leq d,$$

this, together with the fact that $|x_k| \leq R$, $1 \leq k \leq n$, and the definition of R in (4.13), implies that

$$C_4 a_{ii} \leq x_i^2 \leq C_4' a_{ii},\ 1 \leq i \leq d, \qquad (4.21)$$

where C_4 and C_4' are positive constants dependent on a_{ij}, $1 \leq j \leq n$, and $j \neq i$, and $a_{11}, ..., a_{dd}$ are sufficiently large.

Consider the unit cone

$$K_1(\mathbb{R}^d, \mathbb{R}^{n-d}) = \Bigg\{ \mathbf{v} = \begin{pmatrix} v_0 \\ u_0 \end{pmatrix} :\ v_0 = (v_{0,1}, ..., v_{0,d})^T \in \mathbb{R}^d,$$

$$u_0 = (u_{0,1}, ..., u_{0,n-d})^T \in \mathbb{R}^{n-d},\ \text{and } |u_0| \leq |v_0| \Bigg\},$$

where $|v_0| = \max\{|v_{0,1}|, ..., |v_{0,d}|\}$, $|u_0| = \max\{|u_{0,1}|, ..., |u_{0,d}|\}$, and $|\mathbf{v}| = \max\{|v_0|, |u_0|\}$.

Without loss of generality, suppose that $|v_0| = |v_{0,1}|$ and $|u_0| = |u_{0,1}|$. Set $DF(\mathbf{v}) = \begin{pmatrix} v_1 \\ u_1 \end{pmatrix}$, where $v_1 = (v_{1,1}, ..., v_{1,d})^T \in \mathbb{R}^d$, $u_1 = (u_{1,1}, ..., u_{1,n-d})^T \in \mathbb{R}^{n-d}$. So, if a_{11} is sufficiently large, by (4.21), then

$$\begin{aligned} |v_1| \geq |v_{1,1}| &\geq 2|x_1||v_{0,1}| - \sum_{2\leq j\leq d} |a_{1j}||v_{0,j}| - \sum_{d+1\leq j\leq n} |a_{1j}||u_{0,j-d}| \\ &\geq 2|x_1||v_{0,1}| - \sum_{2\leq j\leq n} |a_{1j}||v_{0,1}| \geq \lambda \max_{d+1\leq i\leq n}\Bigg(\sum_{1\leq j\leq n,\ j\neq i} |a_{ij}|\Bigg)|v_{0,1}| \geq \lambda|u_1|, \end{aligned}$$

implying that $m_{\mathcal{C},w_0} \geq \lambda > 1$. This completes the proof. □

Lemma 4.5. In Cases (1)–(3), for fixed parameters a_{ij}, $1 \leq i \neq j \leq n$, and sufficiently large a_{11},...,a_{dd}, the invariant set $\Lambda = \cap_{i=-\infty}^{\infty} F^i(U)$ is uniformly hyperbolic.

Proof. This can be obtained by applying Lemmas 2.1, 4.1–4.4. □

Lemma 4.6. In Cases (1)–(3), for fixed a_{ij}, $1 \leq i \neq j \leq n$, if $a_{11}, ..., a_{dd}$ are sufficiently large, then $F(U) \cap U$ and $F^{-1}(U) \cap U$ have 2^d connected components, respectively.

Proof. It is to show that $F(U) \cap U$ has 2^d components.

Fix any i, $1 \leq i \leq d$, and fix any x_j with $x_j \in [-R, R]$ and $j \neq i$, where R is specified in (4.13). So, the function $f_i(x_1, ..., x_n) = a_{ii} - x_i^2 + \sum_{1\leq j\leq n,\ j\neq i} a_{ij}x_j$ can be rewritten as follows:

$$f_i(x_1, ..., x_{i-1}, y, x_{i+1}, ..., x_n) = a_{ii} - y^2 + \sum_{1\leq j\leq n,\ j\neq i} a_{ij}x_j,\ y \in [-R, R],\ 1 \leq i \leq d.$$

By the definition of R in (4.13), if a_{ii} is sufficiently large, then $f_i(..., y, ...) \approx a_{ii} - y^2 + Ca_{ii}^{1/2} \approx a_{ii} - y^2$, which implies $\mathrm{Graph}(y, f_i(..., y, ...)) \cap ([-R, R] \times [-R, R])$ has two connected components for sufficiently large a_{ii}. Therefore, one has that $F(U) \cap U$ has 2^d components for sufficiently large a_{ii}, $1 \leq i \leq d$.

Since the map F is diffeomorphic, one has that $F^{-1}(U) \cap U = F^{-1}(F(U) \cap U)$ has 2^d components. This completes the proof. □

Therefore, it follows from Lemmas 2.2, 4.5, and 4.6 that Theorem 4.1 holds.

Remark 4.1. Similar results can be obtained, if we substitute the functions $\Delta_i(x)$ by general polynomials. For example, we could consider the following type of polynomial maps:

$$\begin{cases} f_1(x_1,...,x_n) = a_{11}x_1(1-x_1) + \sum_{2\leq j\leq n} a_{1j}x_j \\ \vdots \qquad\quad \vdots \qquad\quad \vdots \qquad\quad \vdots \\ f_d(x_1,...,x_n) = a_{dd}x_d(1-x_d) + \sum_{1\leq j\leq n,\ j\neq d} a_{dj}x_j \\ f_{d+1}(x_1,...,x_n) = \sum_{1\leq j\leq n,\ j\neq d+1} a_{d+1,j}x_j \\ \vdots \qquad\quad \vdots \qquad\quad \vdots \qquad\quad \vdots \\ f_n(x_1,...,x_n) = \sum_{1\leq j\leq n-1} a_{nj}x_j \end{cases}, \tag{4.22}$$

where a_{ij}, $1 \leq i, j \leq n$, are parameters.

Remark 4.2. It is an interesting question to find the general inverse maps for the following type of maps:

$$\begin{cases} f_1(x_1,...,x_n) = a_{11}(x_1-b_{11})\cdots(x_1-b_{1s_1}) + \sum_{2\leq j\leq n} a_{1j}x_j \\ \vdots \qquad\quad \vdots \qquad\quad \vdots \qquad\quad \vdots \\ f_d(x_1,...,x_n) = a_{dd}(x_d-b_{d1})\cdots(x_d-b_{ds_d}) + \sum_{1\leq j\leq n,\ j\neq d} a_{dj}x_j \\ f_{d+1}(x_1,...,x_n) = \sum_{1\leq j\leq n,\ j\neq d+1} a_{d+1,j}x_j \\ \vdots \qquad\quad \vdots \qquad\quad \vdots \qquad\quad \vdots \\ f_n(x_1,...,x_n) = \sum_{1\leq j\leq n-1} a_{nj}x_j \end{cases}, \tag{4.23}$$

where a_{ij} and b_{kl} are parameters. We guess that we could apply similar discussions above to show that for fixed a_{ij}, $1 \leq i \neq j \leq n$ and sufficiently large $a_{11}, ..., a_{dd}$, there might exist a Smale horseshoe and a uniformly hyperbolic invariant set on which the map is topologically conjugate with the fullshift on $s_1s_2\cdots s_d$ symbols.

5 Existence of strange attractors by simulations

In this section, many interesting maps with strange attractors are collected. The Mathematical software Maple is applied to find the strange attractors and the corresponding maximal Lyapunov exponent is calculated. The maximal Lyapunov exponents of these maps are all positive, yielding that these maps have complicated dynamical behavior. For the calculation of the maximal Lyapunov exponents, please refer to [15]. For the simulation of these strange attractors, the initial value is taken as $(0.5, 0.5, 0.5)$, the first 80000 points are omitted and the next 200000 points are kept. Figures 1-6 give the simulation graphs of the maps (2), (8), (11), (20), (24), and (33) from the following table.

Strange Attractors			
No.	Parameters	Polynomial	Max Lya Exp
1.	$a_1 = 2.6$, $a_2 = 0.3$, $a_3 = 0$, $b_1 = 0.4$, $b_2 = 1.7$, $b_3 = 0.5$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	p(x)= x(1-x) q(y)= y(1-y) r(z)=0	0.109706
2.	$a_1 = 2.8$, $a_2 = 0.2$, $a_3 = 0.6$, $b_1 = 0.6$, $b_2 = 2.4$, $b_3 = 0.1$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	p(x)=x(1-x) q(y)=y(1-y) r(z)=0	0.322432
3.	$a_1 = 4.5$, $a_2 = 0.1$, $a_3 = 0$, $b_1 = 0.3$, $b_2 = 7$, $b_3 = 0.1$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x(1-x)^2$ $q(y) = y(1-y)^2$ $r(z) = 0$	0.399491
4.	$a_1 = 5.9$, $a_2 = 0.1$, $a_3 = 0$, $b_1 = 0.3$, $b_2 = 1.3$, $b_3 = 0.1$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x^2(1-x)$ $q(y) = y^2(1-y)$ r(z)=0	0.418139
5.	$a_1 = 5.9$, $a_2 = 0.1$, $a_3 = 0$, $b_1 = 0.3$, $b_2 = 4.7$, $b_3 = 0.1$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x^2(1-x)$ $q(y) = y^2(1-y)$ $r(z) = 0$	0.476862
6.	$a_1 = 8$, $a_2 = 0.1$, $a_3 = 0$, $b_1 = 0.3$, $b_2 = 1.3$, $b_3 = 0.7$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x(1-x)^3$ $q(y) = y(1-y)^3$ $r(z) = 0$	0.312734
7.	$a_1 = 13$, $a_2 = 0.1$, $a_3 = 0$, $b_1 = 0.3$, $b_2 = 8$, $b_3 = 0.1$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x^2(1-x)^2$ $q(y) = y^2(1-y)^2$ $r(z) = 0$	0.427413
8.	$a_1 = 10$, $a_2 = 0.1$, $a_3 = 0$, $b_1 = 0.7$, $b_2 = 10$, $b_3 = 0.2$ $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x(1-x)^4$ $q(y) = y(1-y)^4$ $r(z) = 0$	0.304944
9.	$a_1 = 28$, $a_2 = 0.1$, $a_3 = 0$, $b_1 = 0.3$, $b_2 = 20$, $b_3 = 0.1$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x^2(1-x)^3$ $q(y) = y^2(1-y)^3$ $r(z) = 0$	0.383889
10.	$a_1 = 20$, $a_2 = 0.4$, $a_3 = 0$, $b_1 = 0.3$, $b_2 = 20$, $b_3 = 0.1$ $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x^3(1-x)^2$ $q(y) = y^3(1-y)^2$ $r(z) = 0$	0.434574
11.	$a_1 = 34.5$, $a_2 = 0.4$, $a_3 = 0$, $b_1 = 0.5$, $b_2 = 34$, $b_3 = 0.2$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x^2(1-x)^4$ $q(y) = y^2(1-y)^4$ $r(z) = 0$	0.280940
12	$a_1 = 30$, $a_2 = 0.1$, $a_3 = 0$, $b_1 = 0.2$, $b_2 = 23$, $b_3 = 0.2$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x^2(1-x)^4$ $q(y) = y^2(1-y)^4$ $r(z) = 0$	0.330790
13.	$a_1 = 37$, $a_2 = 0.2$, $a_3 = 0$ $b_1 = 0.6$, $b_2 = 36$, $b_3 = 0.2$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x^3(1-x)^3$, $q(y) = y^3(1-y)^3$ $r(z) = 0$	0.231571
	$a_1 = 29$, $a_2 = 0.2$, $a_3 = 0$,	$p(x) = x^4(1-x)^2$	

14.	$b_1 = 0.6$, $b_2 = 22$, $b_3 = 0.2$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$q(y) = y^4(1-y)^2$ $r(z) = 0$	0.367095
15.	$a_1 = 65$, $a_2 = 0.3$, $a_3 = 0$, $b_1 = 0.5$, $b_2 = 50$, $b_3 = 0.3$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x^3(1-x)^4$ $q(y) = y^3(1-y)^4$ $r(z) = 0$	0.252414
16.	$a_1 = 180$, $a_2 = 0.3$, $a_3 = 0$, $b_1 = 0.2$, $b_2 = 160$, $b_3 = 0.3$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x^4(1-x)^4$ $q(y) = y^4(1-y)^4$ $r(z) = 0$	0.399516
17.	$a_1 = 390$, $a_2 = 0.3$, $a_3 = 0$, $b_1 = 0.5$, $b_2 = 250$, $b_3 = 0.1$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x^4(1-x)^5$ $q(y) = y^4(1-y)^5$ $r(z) = 0$	0.358692
18.	$a_1 = 19000$, $a_2 = 0.1$, $a_3 = 0$, $b_1 = 0.3$, $b_2 = 800$, $b_3 = 0.7$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = x^8(1-x)^7$ $q(y) = y^6(1-y)^7$ $r(z) = 0$	0.542847
19.	$a_1 = 2.5$, $a_2 = 0.2$, $a_3 = 0$, $b_1 = 0.2$, $b_2 = 2.7$, $b_3 = 0.1$, $c_1 = 1$, $c_2 = 0$, $c_3 = 0$	$p(x) = (x+1)x(1-x)$ $q(y) = (y+1)y(1-y)$ $r(z) = 0$	0.210240
20.	$a_1 = 0$, $a_2 = 1$, $a_3 = 0$, $b_1 = 0$, $b_2 = 0$, $b_3 = 1$, $c_1 = 0.6$, $c_2 = 0.6$, $c_3 = 1$	$p(x) = 0$ $q(y) = 0$ $r(z) = (z+1)^2(1-z) - 1$	0.175343
21.	$a_1 = 0$, $a_2 = 1$, $a_3 = 0$, $b_1 = 0$, $b_2 = 0$, $b_3 = 1$, $c_1 = 0.1$, $c_2 = 0.2$, $c_3 = 14.5$	$p(x) = 0$ $q(y) = 0$ $r(z) = z^2(1-z)^2$	0.401407
22.	$a_1 = 0$, $a_2 = 1$, $a_3 = 0$, $b_1 = 0$, $b_2 = 0$, $b_3 = 1$, $c_1 = 0.3$, $c_2 = 0.4$, $c_3 = 14.1$	$p(x) = 0$ $q(y) = 0$ $r(z) = z^3(1-z)^2$	0.250262
23.	$a_1 = 0$, $a_2 = 0.7$, $a_3 = 0$, $b_1 = 0$, $b_2 = 0$, $b_3 = 0.7$, $c_1 = 0.2$, $c_2 = 0$, $c_3 = 23$	$p(x) = 0$ $q(y) = 0$ $r(z) = z^3(1-z)^2$	0.492135
24.	$a_1 = 0$, $a_2 = 0.8$, $a_3 = 0$, $b_1 = 0$, $b_2 = 0$, $b_3 = 0.7$, $c_1 = 0.2$, $c_2 = 0$, $c_3 = 53$	$p(x) = 0$ $q(y) = 0$ $r(z) = z^2(1-z)^4$	0.510269
25.	$a_1 = 0$, $a_2 = 0.8$, $a_3 = 0$, $b_1 = 0$, $b_2 = 0$, $b_3 = 0.7$, $c_1 = 0.3$, $c_2 = 0.3$, $c_3 = 440$	$p(x) = 0$ $q(y) = 0$ $r(z) = z^4(1-z)^5$	0.279378
26.	$a_1 = 2$, $a_2 = 0.3$, $a_3 = 0$, $b_1 = 0.4$, $b_2 = 2$, $b_3 = 0.5$, $c_1 = 0.5$, $c_2 = 0$, $c_3 = 2$	$p(x) = x(1-x)$ $q(y) = y(1-y)$ $r(z) = z(1-z)$	0.113261
27.	$a_1 = 3.5$, $a_2 = 0.1$, $a_3 = 0$, $b_1 = 0.3$, $b_2 = 2.3$, $b_3 = 0.2$, $c_1 = 0.2$, $c_2 = 0$, $c_3 = 3$	$p(x) = x(1-x)$ $q(y) = y(1-y)$ $r(z) = z(1-z)$	0.336029
28.	$a_1 = 3.2$, $a_2 = 0.1$, $a_3 = 0$, $b_1 = 0.3$, $b_2 = 2.5$, $b_3 = 0.2$,	$p(x) = x(1-x)$ $q(y) = y(1-y)$	0.0399768

	$c_1 = 0.2,\ c_2 = 0,\ c_3 = 2.6$	$r(z) = z(1-z)$	
29.	$a_1 = 3.2,\ a_2 = 0.1,\ a_3 = 0,$ $b_1 = 0.3,\ b_2 = 2.5,\ b_3 = 0.2,$ $c_1 = 0.2,\ c_2 = 0,\ c_3 = 3$	$p(x) = x(1-x)$ $q(y) = y(1-y)$ $r(z) = z(1-z)$	0.0971938
30.	$a_1 = 3.2,\ a_2 = 0.1,\ a_3 = 0,$ $b_1 = 0.3,\ b_2 = 2.5,\ b_3 = 0.2,$ $c_1 = 0.2,\ c_2 = 0,\ c_3 = 3.2$	$p(x) = x(1-x)$ $q(y) = y(1-y)$ $r(z) = z(1-z)$	0.198463
31.	$a_1 = 3.2,\ a_2 = 0.1,\ a_3 = 0,$ $b_1 = 0.3,\ b_2 = 2.5,\ b_3 = 0.2,$ $c_1 = 0.2,\ c_2 = 0,\ c_3 = 3.3$	$p(x) = x(1-x)$ $q(y) = y(1-y)$ $r(z) = z(1-z)$	0.226691
32.	$a_1 = 3.2,\ a_2 = 0.1,\ a_3 = 0,$ $b_1 = 0.3,\ b_2 = 2.5,\ b_3 = 0.2,$ $c_1 = 0.2,\ c_2 = 0,\ c_3 = 3.4$	$p(x) = x(1-x)$ $q(y) = y(1-y)$ $r(z) = z(1-z)$	0.293952
33.	$a_1 = 15,\ a_2 = 0.2,\ a_3 = 0.1,$ $b_1 = 0.3,\ b_2 = 2.6,\ b_3 = 0.2,$ $c_1 = 0.6,\ c_2 = 0.2,\ c_3 = 20$	$p(x) = x^2(1-x)^4$ $q(y) = y(1-y)$ $r(z) = z^2(1-z)^3$	0.149243
34.	$a_1 = 20,\ a_2 = 0.3,\ a_3 = 0.2,$ $b_1 = 0.2,\ b_2 = 2.5,\ b_3 = 0.2,$ $c_1 = 0.2,\ c_2 = 0.2,\ c_3 = 25$	$p(x) = x^2(1-x)^4$ $q(y) = y(1-y)$ $r(z) = z^2(1-z)^3$	0.183814
35.	$a_1 = 4,\ a_2 = 0.4,\ a_3 = 0.3,$ $b_1 = 0.2,\ b_2 = 90,\ b_3 = 0.2,$ $c_1 = 0.2,\ c_2 = 0.2,\ c_3 = 1760$	$p(x) = x^2(1-x)^3$ $q(y) = y^3(1-y)^4$ $r(z) = z^5(1-z)^6$	0.333486
36.	$a_1 = 4,\ a_2 = 0.4,\ a_3 = 0.3,$ $b_1 = 0.2,\ b_2 = 60,\ b_3 = 0.2,$ $c_1 = 0.2,\ c_2 = 0.2,\ c_3 = 1700$	$p(x) = x^2(1-x)^3$ $q(y) = y^3(1-y)^4$ $r(z) = z^5(1-z)^6$	0.206663

6 Example

In this section, two examples are provided to illustrate the theoretical results obtained in Theorems 3.2 and 3.5. All the illustration graphs in this section are drawn by using the software Mathematica. The interested readers can make simple programs to run some softwares and obtain more interesting graphs.

Example 6.1. Consider the following example:

$$\begin{cases} f_1(x,y,z) = 7(x-1)(2-x) + 0.2y + z \\ f_2(x,y,z) = x \\ f_3(x,y,z) = 0.2y, \end{cases} \tag{6.1}$$

It is evident that this map satisfies the conditions in Case (i) of Theorem 3.2. Set $U = [1,2] \times [1,2] \times [0.2, 0.4]$. Figures 7 and 8 provide the graphs of $F(U) \cap U$ and $F^{-1}(U) \cap U$, respectively. By Theorem 3.2, there exists a Smale horseshoe, the invariant set is uniformly hyperbolic and is topologically conjugate to fullshift on two symbols.

Example 6.2. Consider the following example:

$$\begin{cases} f_1(x,y,z) = 7(x-1)(2-x) + 0.4y \\ f_2(x,y,z) = 7(y-1)(2-y) + 0.4z \\ f_3(x,y,z) = x, \end{cases} \tag{6.2}$$

It is evident that this map satisfies the conditions in Case (i) of Theorem 3.5. Set $U = [1,2] \times [1,2] \times [1,2]$. Figures 9 and 10 give the graphs of $F(U) \cap U$ and $F^{-1}(U) \cap U$, respectively. From the computer simulations, we find that the graph of $F^{-1}(U)$ is very big compared with U. So, we only give a part of the graph of $F^{-1}(U) \cap U$. It follows from Theorem 3.5 that there are a Smale horseshoe and the uniformly hyperbolic invariant set on which the map is topologically conjugate to fullshift on four symbols.

Acknowledgments

I would like to thank Professor Sheldon Newhouse for his encouragement, comments, and providing many useful references.

I devote this work to my parents, I cannot thank my parents enough for all the support and love they have given me.

References

[1] J. Banks, J. Brooks, G. Cairns, G. Davis, and P. Stacey. On devaney's definition of chaos. *Amer. Math. Mon.*, 99:332–334, 1992.

[2] H. Bass, E. Connell, and D. Wright. The jacobian conjecture: reduction of degree and formal expansion of the inverse. *Bull. Amer. Soc.*, 7:287–330, 1982.

[3] E. Bedford and J. Smillie. Polynomial diffeomorphisms of $\mathbf{C}^2$. vi: Connectivity of j. *Ann. of Math.*, 148:695–735, 1998.

[4] E. Bedford and J. Smillie. Polynomial diffeomorphisms of $\mathbf{C}^2$. vii: Hyperbolicity and external rays. *Ann. Sci. Ecole Norm. Sup. 4 série*, 32:455–497, 1999.

[5] E. Bedford and J. Smillie. Polynomial diffeomorphisms of $\mathbf{C}^2$. viii: Quasiexpansion. *Amer. Journal of Math.*, 124:221–271, 2002.

[6] E. Bedford and J. Smillie. Real polynomial diffeomorphisms with maximal entropy: I. tangencies. *Ann. of Math.*, 160:1–26, 2004.

[7] E. Bedford and J. Smillie. Real polynomial diffeomorphisms with maximal entropy: Ii. small jacobian. *Ergod. Th. & Dynam. Sys.*, 26:1259–1283, 2005.

[8] F. Belkhouche and U. Qidwai. Binary image encoding using 1d chaotic maps. *IEEE Region 5, Ann. Techn. Conf.*, pages 39–43, 2003.

[9] M. Benedicks and L. Carleson. The dynamics of the hénon map. *Ann. of Math.*, 133:73–169, 1991.

[10] M. Benedicks and M. Viana. Solution of the basin problem for hénon-like attractors. *Invent. Math.*, 143:375–434, 2001.

[11] M. Benedicks and L. S. Young. Sinai-bowen-ruelle measure for certain hénon maps. *Invent. Math.*, 112:541–576, 1993.

[12] Y. Cao, S. Luzzato, and I. Rios. The boundary of hyperboliity for hénon-like families. *Ergod. Th. & Dynam. Sys.*, 28:1049–1080, 2008.

[13] G. Chen and X. Dong. On feedback control of chaotic continuous-time systems. *IEEE Trans. Circ. Syst.-I*, 40:591–601, 1993.

[14] C. Cheng, S. Wang, and J. Yu. Degree bounds for inverses of polynomial automorphisms. *Proc. Amer. Math. Soc.*, 120:705–707, 1994.

[15] G. H. Choe. *Computational Ergodic Theory.* Springer, Berlin Heidelberg, 2005.

[16] R. L. Devaney. *An Introduction to Chaotic Dynamical Systems, 2nd ed.* Addison-Wesley, New York, 1989.

[17] R. L. Devaney and Z. Nitecki. Shift automorphisms in the hénon mapping. *Commun. Math. Phys.*, 67:137–146, 1979.

[18] H. R. Dullin and J. D. Meiss. Generalized hénon maps: the cubic diffeomorphisms of the plane. *Phys. D*, 143:262–289, 2000.

[19] J. P. Eckmann and D. Ruelle. Ergodic theory of chaos and strange attractors. *Rev. Mod. Phys.*, 57:617–656, 1985.

[20] Z. Elhadj and J. C. Sprott. Classification of three-dimensional quadratic diffeomorphisms with constant jacobian. *Front. Phys. China*, 4:111–121, 2009.

[21] J. Feng and C. K. Tse. *Reconstruction of Chaotic Signals with Applications to Chaos-based Communications.* World Scientific, SingaporeCTsinghua University Press, Beijing, 2007.

[22] D. Fournier-Prunaret, R. Lopez-Ruiz, and A. K. Taha. Route to chaos in three-dimensional maps of logistic type. *Grazer Mathematische Berichte*, 350:82–95, 2006.

[23] S. Friedland and J. Milnor. Dynamical properties of plane polynomial automorphisms. *Ergod. Th. & Dynam. Sys.*, 9:67–99, 1989.

[24] S. V. Gonchenko and M. C. Li. Towards a hyperbolic dynamics of three-dimensional hénon-like maps, part i. *preprint*, 2008.

[25] S. V. Gonchenko, J. D. Meiss, and I. I. Ovsyannikov. Chaotic dynamics of three-dimensional hénon maps that originate from a homoclinic bifurcation. *Regular and Chaotic Dynamics*, 11:191–212, 2006.

[26] S. V. Gonchenko, I. I. Ovsyannikov, C. Simó, and D. Turaev. Three-dimensional hénon-like maps and wild lorenz-like attractors. *Int. J. Bifurcation and Chaos*, 15:3493–3508, 2005.

[27] S. V. Gonchenko, L. P. Shilnikov, and D. V. Turaev. On global bifurcations in three-dimensional diffeomorphisms leading to wild lorenz-like attractors. *Regular and Chaotic Dynamics*, 14:137–147, 2009.

[28] M. Hénon. A two-dimensional mapping with a strange attractor. *Commun. Math. Phys.*, 50:69–77, 1976.

[29] M. Jessa. Data encryption algorithms using onedimensional chaotic maps. *Circuits and Systems. Proc. IEEE Int. Symp. Circuits and Systems, Geneva*, 1:711–714, 2000.

[30] D. Kelsey. The economics of chaos or the chaos of economics. *Oxford Econ. Papers*, 40:1–31, 1988.

[31] T. Kohda, Y. Ookubo, and K. Ishii. A color image communication using yiq signals by spread spectrum techniques. *Proc. IEEE Int. Symp. Spread Spectrum Techn. Appl.*, 3:743–747, 1998.

[32] Q. Li and X. Yang. A 3d smale horseshoe in a hyperchaotic discrete-time system. *Discrete Dynamics in Nature and Society*, 2007:Article ID 16239, 9 pages, 2007.

[33] T. Li and J. A. Yorke. Period three implies chaos. *Am. Math. Mon.*, 82:985–992, 1975.

[34] E. Lorenz. Deterministic non-periodic flow. *J. Atmos. Sci.*, 20:130–141, 1963.

[35] R. May. Simple mathematical models with very complicated dynamics. *Nature*, 261:45–67, 1976.

[36] T.-T. Moh. On the jacobian conjecture and the configuration of roots. *J. Reine Angew. Math.*, 340:140–212, 1983.

[37] J. Moser. On quadratic symplectic mappings. *Math. Z.*, 216:417–430, 1994.

[38] N. Murray and M. Holman. The origin of chaotic in the outer solar system. *Science*, 283:1877–1881, 1999.

[39] S. Newhouse. *Cone-fields, domination, and hyperbolicity*, pages 419–432. Modern Dynamical Systems and Applications. Cambridge Univ. Press, Cambridge, 2004.

[40] H. J. Poincaré. Sur le problème des trois corps et les équations de la dynamique. *Acta Mathematica*, 13:1–270, 1890.

[41] H. J. Poincaré. *Les Méthodes Nouvelles de la Mécanique Celeste, Vols. 1–3.* Gauthiers-Villars,Paris, 1892, 1893, 1899. English translation edited by D. Goroff, Amer. Institute of Physics, New York, 1993.

[42] R. G. L. Pryor and J. E. H. Bright. Applying chaos theory to careers: attraction and attractors. *J. Vocat. Behav.*, 71:375–400, 2007.

[43] C. Robinson. *Dynamical Systems: Stability, Symbolic Dynamics and Chaos.* CRC Press, Florida, 1999.

[44] Y. Shi, H. Ju, and G. Chen. Coupled-expanding maps and one-sided symbolic dynamical systems. *Chaos Solit. Fract.*, 39:2138–2149, 2009.

[45] C. Sparrow. *The Lorenz Equations: Bifurcations, Chaos, and Strange Attractors.* Springer-Verlag, NY, 1982.

[46] J. C. Sprott. *Strange Attractors: Creating Patterns in Chaos.* M&T Books, NY, 1993.

[47] J. C. Sprott. Can a monkey with a computer create art? *Nonlinear Dynamics, Psychology, and Life Sciences*, 8:103–114, 2004.

[48] P. E. Strizhak. Application of chemical chaos to analytical chemistry. *Adv. Compl. Syst.*, 6:137–153, 2003.

[49] A. Tsuneda. On auto-correlation properties of chaotic binary sequences generated by onedimensional maps. *Proc. Industr. Electron. Soc. Ann. Conf.*, 3:2025–2030, 2000.

[50] S. Wang. A jacobian criterion for separability. *J. Algebra*, 65:453–494, 1980.

[51] S. Wiggins. *Introduction to Applied Nonlinear Dynamical System and Chaos.* Springer-Verlag, 1990.

[52] X. Zhang. Hyperbolic invariant sets of the real generalized hénon maps. *Chaos, Solitons, Fractals*, 43:31–41, 2010.

[53] X. Zhang. Hyperbolicity of the invariant sets for the real polynomial maps. *Chaos, Solitons, Fractals*, 45:314–324, 2012.

[54] X. Zhang and Y. Shi. Coupled-expanding maps for irreducible transiton matrices. *Int. J. Bifurcation and Chaos*, 20:3769–3783, 2010.

[55] X. Zhang, Y. Shi, and G. Chen. Some properties of coupled-expanding maps in compact sets. *Proc. Amer. Math. Soc.*, 141(2):585–595, 2013.

List of Figure Captions

Figure 1. The chaotic attractor of map (2) in Section 5, where the initial value is taken as $(0.5, 0.5, 0.5)$.

Figure 2. The chaotic attractor of system map (8) in Section 5, where the initial value is taken as $(0.5, 0.5, 0.5)$.

Figure 3. The chaotic attractor of map (11) in Section 5, where the initial value is taken as $(0.5, 0.5, 0.5)$.

Figure 4. The chaotic attractor of map (20) in Section 5, where the initial value is taken as $(0.5, 0.5, 0.5)$.

Figure 5. The chaotic attractor of map (24) in Section 5, where the initial value is taken as $(0.5, 0.5, 0.5)$.

Figure 6. The chaotic attractor of map (33) in Section 5, where the initial value is taken as $(0.5, 0.5, 0.5)$.

Figure 7. The illustration graph of $F(U) \cap U$ in Example 6.1, where $U = [1, 2] \times [1, 2] \times [0.2, 0.4]$.

Figure 8. The illustration graph of $F^{-1}(U) \cap U$ in Example 6.1, where $U = [1, 2] \times [1, 2] \times [0.2, 0.4]$.

Figure 9. The illustration graph of $F(U) \cap U$ in Example 6.2, where $U = [1, 2] \times [1, 2] \times [1, 2]$.

Figure 10. The illustration graph of $F^{-1}(U) \cap U$ in Example 6.2, where $U = [1, 2] \times [1, 2] \times [1, 2]$.



Figure 1: The chaotic attractor of map (2) in Section 5, where the initial value is taken as $(0.5, 0.5, 0.5)$.



Figure 2: The chaotic attractor of system map (8) in Section 5, where the initial value is taken as $(0.5, 0.5, 0.5)$.



Figure 3: The chaotic attractor of map (11) in Section 5, where the initial value is taken as $(0.5, 0.5, 0.5)$.



Figure 4: The chaotic attractor of map (20) in Section 5, where the initial value is taken as $(0.5, 0.5, 0.5)$.



Figure 5: The chaotic attractor of map (24) in Section 5, where the initial value is taken as $(0.5, 0.5, 0.5)$.



Figure 6: The chaotic attractor of map (33) in Section 5, where the initial value is taken as $(0.5, 0.5, 0.5)$.



Figure 7: The illustration graph of $F(U) \cap U$ in Example 6.1, where $U = [1,2] \times [1,2] \times [0.2, 0.4]$.



Figure 8: The illustration graph of $F^{-1}(U) \cap U$ in Example 6.1, where $U = [1,2] \times [1,2] \times [0.2, 0.4]$.



Figure 9: The illustration graph of $F(U) \cap U$ in Example 6.2, where $U = [1,2] \times [1,2] \times [1,2]$.



Figure 10: The illustration graph of $F^{-1}(U) \cap U$ in Example 6.2, where $U = [1,2] \times [1,2] \times [1,2]$.